arXiv:1204.5308v1 [math.GT] 24 Apr 2012

THE TOPOLOGICAL AND GEOMETRICAL FINITENESS OF COMPLETE FLAT LORENTZIAN 3-MANIFOLDS WITH FREE FUNDAMENTAL GROUPS

SUHYOUNG CHOI

Abstract. We prove the topological tameness of a 3-manifold with a free fundamental group admitting a complete flat Lorentzian metric; i.e., a Margulis space-time isomorphic to the quotient of the complete flat Lorentzian space by the free and properly discontinuous isometric action of the free group of rank ≥ 2. We will use our particular point of view that a Margulis space-time is a real projective manifold in an essential way. The basic tools are a bordification by a closed real projective surface with a free holonomy group, the important work of Goldman, Labourie, and Margulis on geodesics in the Margulis space-times and the 3-manifold topology. Finally, we show that Margulis space-times are geometrically finite under our definition.

Our work goes back to the works of mathematicians on complete affine manifolds; i.e., the manifolds that are quotient spaces of the affine space by the properly discontinuous and free actions of groups of affine transformations. Auslander conjectured that a compact complete affine manifold has a solvable fundamental group. Goldman, Fried, and Hirsch [22] and [21] proved this conjecture for three-dimensional complete compact affine manifolds. This conjecture was generalized by Milnor who asked whether a complete affine manifold has a solvable fundamental group. However, Margulis [32] found a counter-example by considering a Lorentzian 3-space $\mathsf{E}^{2,1}$ with a free group of rank two acting on it by deforming a linear Lorentzian isometry group. The quotient space is now said to be a Margulis space-time. Drumm [15] found fundamental domains of these actions. Charette, Drumm, and Goldman [7] classified these spaces for some classes of examples and found fundamental domains of these examples using crooked planes. Goldman, Labourie, and Margulis [28] and [27] constructed generalizations of Margulis invariants and they characterized the deformation spaces of Margulis space-times in terms of the invariants.

Date: April 24, 2012.

2000 *Mathematics Subject Classification.* Primary 57M50; Secondary 53C15, 53C50, 53A20.

Key words and phrases. Lorentzian manifolds, geometric structure, geometric finiteness.

this work was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MEST) (No.2010-0027001).

A question of Goldman was whether these spaces were topologically tame. (See [17] and [7].) We will answer this in the positive. However, when the Margulis space-time has parabolics, this question is still open. The Auslander conjecture still remains open for dimension $n \geq 7$.

The theory of compactifying open manifolds goes back to Brouwder, Levine and Livsay [6] and Siebenmann [37]. For 3-manifolds, Tucker [40] and Scott and Tucker [39] made initial clarification. See also Meyers [34]. As a note, we state that complete hyperbolic 3-manifolds with finitely generated fundamental groups were shown to be tame by Agol, Calegari, and Gabai. See Bowditch [5] for details. We won't elaborate on this as the methods are completely different. Our proof is more close to the proof of the tameness of geometrically finite hyperbolic 3-manifolds due to Thurston [38].

Let $\mathsf{V}^{2,1}$ denote the vector space $\mathbb{R}^3$ with a Lorentzian norm of sign $1, 1, -1$, and the Lorentzian space-time $\mathsf{E}^{2,1}$ can be thought of as the vector space with translation by any vector allowed. A *Lorentzian isometry* is an automorphism of the space preserving the Lorentzian norm and the group of Lorentzian isometries is denoted by $\mathsf{Isom}(\mathsf{E}^{2,1})$. We will concerned ourselves with only the subgroup $\mathsf{Isom}^+(\mathsf{E}^{2,1})$ of orientation-preserving isometries.

The projectivization $\mathsf{P}(\mathsf{V}^{2,1} - \{O\})$ is defined as the quotient space

$$\mathsf{V}^{2,1} - \{O\}/\sim \text{ where } \mathsf{v} \sim \mathsf{w} \text{ if and only if } \mathsf{v} = s\mathsf{w} \text{ for } s \in \mathbb{R} - \{0\}.$$

Of course, $\mathsf{P}(\mathsf{V}^{2,1} - \{O\})$ is identical with the real projective 2-space $\mathbb{R}\mathsf{P}^2$. Recall that we imbed the hyperboloid model $\mathbb{H}^2$ of the hyperbolic plane to the disk that is the interior of conic in $\mathbb{R}P^2$ determined by null vectors. Then the image $\mathbb{S}_+$ of the space of future time-like vectors can be identified with the hyperbolic 2-plane $\mathbb{H}^2$, which is basically the Beltrami-Klein model of the hyperbolic plane. The linear group $\mathsf{SO}(2,1)$ acts faithfully on $\mathbb{H}^2 = \mathbb{S}_+$ as the orientation-preserving isometry group and acts on $\mathsf{P}(\mathsf{V}^{2,1} - \{O\})$ where the action is induced from that on $\mathsf{V}^{2,1}$.

Recall that there is a homomorphism

$$\mathcal{L} : \mathsf{Isom}^+(\mathsf{E}^{2,1}) \rightarrow \mathsf{SO}(2,1)$$

given by taking the linear part. A Lorentzian transformation is said to be *parabolic* if its linear part has only eigenvalues 1 but is not identity.

Suppose that Γ is a finitely generated Lorentzian isometry group acting freely and properly on $\mathsf{E}^{2,1}$. We assume that Γ is not amenable (i.e., not solvable). Then $\mathsf{E}^{2,1}/\Gamma$ is said to be a *Margulis space-time.* Γ injects under $\mathcal{L}$ to $\mathcal{L}(\Gamma)$ acting properly discontinuously and freely on $\mathbb{S}_+$. By Mess [33], Γ is a free group of rank ≥ 2. (See [22] where this nontrivial fact was first discovered and Section 3 of [27] for details.) Thus, we restrict ourselves to the group actions of free groups of rank ≥ 2 in this paper.

A *real projective structure* on a manifold is given by an atlas of charts to $\mathbb{R}\mathsf{P}^2$ with projective transition maps. Such geometric structures were first considered by Kuiper, Koszul, Benzecri and so on in the 1950s and 1960s.

Further developments can be seen in Goldman [26], Choi and Goldman [12], and Cooper, Long, and Thistlethwaite [13] and [14].

Let us denote by $\mathbb{S}$ the double-cover of $\mathbb{R}\mathsf{P}^2$ where $\mathbb{S}$ will be realized as the space of (oriented) directions in $\mathsf{E}^{2,1}$. Hence, $\mathsf{SO}(2,1)$ acts on $\mathbb{S}$ locally projectively. We will use a closed real projective surface that is the quotient of a domain in $\mathbb{S}$.

Theorem 1. *Let $\Gamma \subset \mathsf{SO}(2,1)$ be a free group of rank ≥ 2 acting on the hyperbolic 2-space $\mathbb{H}^2$ properly discontinuously and freely but without any parabolic holonomy. Let $\mathbb{S} := \mathbb{S}(\mathsf{V}^{2,1})$ be the sphere of directions in $\mathsf{V}^{2,1}$, which is a model for the universal covering space $\widehat{\mathbb{R}\mathsf{P}^2} \longrightarrow \mathbb{R}\mathsf{P}^2$. Then there exists a Γ-invariant open domain $\mathcal{D} \subset \mathbb{S}(\mathsf{V}^{2,1})$ such that $\mathcal{D}/\Gamma$ is a closed surface Σ.*

These surfaces correspond to closed real projective surfaces of genus g, $g \geq 2$, discovered by Goldman [24] in the late 1970s. Goldman did not notice that although their developing maps from the universal cover $\tilde{\Sigma}$ of some such surface Σ are not covering-spaces onto their images in $\mathbb{R}\mathsf{P}^2$, when lifted to the double-covering space $\mathbb{S}(\mathsf{V}^{2,1}) := \widehat{\mathbb{R}\mathsf{P}^2} \longrightarrow \mathbb{R}\mathsf{P}^2$, the developing maps (remarkably) are covering-spaces onto open domains. That is, the surface is a quotient of a domain in $\mathbb{S}$ by a group of projective automorphisms. This is an $\mathbb{R}\mathsf{P}^2$-analog of the standard Schottky uniformization of a Riemann surface as a $\mathbb{C}\mathsf{P}^1$-manifold as observed by Goldman.

An n-dimensional open manifold is said to be *tame* if it is homeomorphic to the interior of a compact n-manifold with boundary. We obtain a *handle body* is a 3-dimensional manifold from a 3-ball B^3 by attaching $\coprod_{i=1}^{k} D_i^2 \times I$ by imbedding $\coprod_{i=1}^{k} D_i^2 \times \partial I$ into ∂B^3 where D_i^2 is a standard disk for each i and I is the unit interval with boundary ∂I.

Theorem 2. *Let M be $\mathsf{E}^{2,1}/\Gamma$ where $\mathsf{E}^{2,1}$ is the standard Lorentzian space with the standard norm given by $x_0^1 - x_1^2 - x_2^2$, and let Γ be a nonamenable orientation-preserving free Lorentzian isometry group acting freely and properly discontinuously on $\mathsf{E}^{2,1}$. Assume that the linear part of Γ does not contain any parabolic elements. Then M is homeomorphic to the interior of a handle body of genus equal to the rank of the free group Γ.*

Originally, we thought of a proof using the constant Gauss curvature surfaces of Labourie years ago. However, we have not been able to complete it for a while. Here, we came up with a simpler approach using compactifications and Dehn's lemma.

We define in this paper that a complete geodesic on a quotient space of a hyperbolic space or a Lorentzian space is *weakly recurrent* if it is bounded in both directions; that is, the closure of the forward part is compact and so is that of the backward part; i.e., the α-limit and the ω-limit are both nonempty and compact. This is the same as the term "recurrent" in [27] and [8], which does not agree with the common usage in the dynamical system theory whereas the recurrence set they discuss is the same as the nonwandering set as in Eberlein [19] and Katok and Hasselblatt [30]: the reason is that these are geodesics with both endpoints in the limit sets and they coincide with the nonwondering set in [19] by Corollary 3.8 there.

In some sense, the work of Goldman, Labourie, and Margulis [28] on weakly recurrent geodesics on M implies tameness: A Margulis space-time is geometrically finite in some sense since every closed geodesic is in a bounded neighborhood. We will explain our particular viewpoints on this in Section six.

In Section one, we review facts on real projective structures and geometry that we need. We compactify $\mathsf{E}^{2,1}$ by a real projective 2-sphere $\mathbb{S}$ that is just the space of directions in $\mathsf{E}^{2,1}$. (The discussion on arcs in Sections 1.5 and 1.6 were written jointly with Goldman.)

In Section two, we prove Theorem 1. This will be done by finding an open domain $\tilde{\Sigma}$ in the boundary $\mathbb{S}$ of $\mathsf{E}^{2,1}$ where Γ acts properly discontinuously and freely. $\tilde{\Sigma}/\Gamma$ is shown to be a closed real projective surface Σ. Such a surface was constructed by Goldman [24]; however, we realize it as the quotient of an open domain in the sphere $\mathbb{S}$, which is a union of two disks and infinitely many strips joining the two.

Now let Γ be as assumed in Theorem 2. In Section three, we recall the work of Goldman, Labourie, and Margulis [28] on weakly recurrent geodesics on the Margulis space-time. Their work in fact shows that all weakly recurrent space-like geodesics in a Margulis space-time are in a uniformly bounded part.

In Section four, we prove the proper discontinuity of the action of the group Γ on $\mathsf{E}^{2,1} \cup \tilde{\Sigma}$. We first show that for a fixed Lorentzian isometry g, we can push every compact convex subset of $\mathsf{E}^{2,1} \cup \tilde{\Sigma}$ into an ϵ-neighborhood of the attracting segment of g in $\mathbb{S}$ by g^i for sufficiently large i. (This is the key model of our proof.)

We next show that for any compact set $K \subset \mathsf{E}^{2,1} \cup \tilde{\Sigma}$, there are only finitely many $g \in \Gamma$ so that $g(K) \cap K \neq \emptyset$. Suppose not. We find a sequence $\{g_i\}$ so that $g_i(K) \cap K \neq \emptyset$. Using the work [28], we show that we can find an infinite sequence $\{g_i\}$ that behave almost like $\{g^i\}$ in the dynamical sense as $i \to \infty$. In other words, we can find a uniformly bounded sequence of coordinates to normalize g_i into fixed forms and these coordinates are convergent.

In Section five, we prove the tameness of $\mathsf{E}^{2,1}/\Gamma$, i.e., Theorem 2, using the classical 3-manifold topology as developed by Dehn and so on.

In Section six, we show that there exists an invariant set of hyperplanes asymptotic to the limit set Λ of Γ on $\mathrm{Cl}(\mathbb{S}_+)$. This gives us two disjoint Γ-invariant closed convex domains in $\mathsf{E}^{2,1}$ asymptotic to Λ and its antipodal set Λ_- in $\mathbb{S}$. (This is a generalization of a result of Mess [33].)

In Section seven, we also define "geometric finiteness" in this setting and show that there exists a topological core of a Margulis space-time that contains all space-like weakly recurrent geodesics.

This work was started as a joint work with William Goldman from his question on the bordification of the Margulis space-time during the Newton Institute Workshop titled "Representations of Surface Groups and Higgs Bundles" held in Oxford during March 14th to 18th. The bordification idea is originally due to Charles Francis [20] where he found a bordifications of the Margulis space-times in the conformal sense.

We benefited from discussions with Virginie Charette, Todd Drumm, and Francois Labourie clarifying the issues here. We also thank Yves Coudene for clarifying issues with geodesic flows. Of course, the author is very much indebted to William Goldman for discussing with him through many phases of this work for which he is very grateful. The author thanks the Newton Institute and the Institut Henri Poincaré where parts of this work were carried out.

1. PRELIMINARY

1.1. Real projective structures on manifolds.

The projective space is given by

$$\mathbb{R}\mathsf{P}^n = (\mathbb{R}^{n+1} - \{O\})/\sim \ \text{ where } \mathsf{v} \sim \mathsf{w} \text{ if } \mathsf{v} = s\mathsf{w} \text{ for } s \in \mathbb{R} - \{0\}.$$

The general linear group is sent

$$\mathsf{GL}(n+1,\mathbb{R}) \to \mathsf{PGL}(n+1,\mathbb{R}) := \mathsf{GL}(n+1,\mathbb{R})/\sim$$

where two linear maps L_1 and L_2 are equivalent if $L_1 = sL_2$ for $s \in \mathbb{R} - \{0\}$. A map of an open subset of $\mathbb{R}\mathsf{P}^n$ to one of $\mathbb{R}\mathsf{P}^n$ is *projective* if it is a restriction of an element of $\mathsf{PGL}(n+1,\mathbb{R})$.

The projective geometry is given by a pair $(\mathbb{R}\mathsf{P}^n, \mathsf{PGL}(n+1,\mathbb{R}))$. A *real projective structure* on a manifold M is given by a maximal atlas of charts to $\mathbb{R}\mathsf{P}^n$ with projective transition maps. The manifold M with a real projective structure is said to be a *real projective manifold.* A *projective map* for two real projective manifolds M and N is a map $f : M \to N$ so that $\phi \circ f \circ \psi^{-1}$ is projective whenever it is defined where ϕ is a chart for N and ψ^{-1} is a local inverse of a chart for M.

A real projective structure on a manifold M gives us an immersion $\mathsf{dev} : \tilde{M} \to \mathbb{R}\mathsf{P}^n$ that is equivariant with respect to a homomorphism $h : \pi_1(M) \to \mathsf{PGL}(n+1,\mathbb{R})$: i.e.,

$$\mathsf{dev} \circ \gamma = h(\gamma) \circ \mathsf{dev} \text{ for every } \gamma \in \pi_1(M).$$

Here, dev is called a *developing map* and h is called the *holonomy homomorphism.* (dev, h) is only determined up to an action of $\mathsf{PGL}(n+1,\mathbb{R})$ where

$$g(\mathsf{dev}, h(\cdot)) = (g \circ \mathsf{dev}, gh(\cdot)g^{-1}) \text{ for } g \in \mathsf{PGL}(n+1,\mathbb{R}).$$

Conversely, such a pair (dev, h) will give us a real projective structure since dev gives us charts that have projective transition maps. (See [10] for details.) Furthermore, given a regular cover $\hat{M}$ of M and an immersion

$\widehat{D} : \hat{M} \to \mathbb{R}\mathsf{P}^n$ equivariant with respect to a homomorphism $\hat{h} : G \to \mathsf{PGL}(n+1, \mathbb{R})$ for the deck transformation group G of $\hat{M}$, we also obtain a real projective structure on M.

We can identify $\mathbf{S}^n$ with

$$(\mathbb{R}^{n+1} - \{O\})/\sim \text{ where } \mathsf{v} \sim \mathsf{w} \text{ if } \mathsf{v} = s\mathsf{w} \text{ for } s > 0.$$

There exists a quotient map $q_1 : \mathbf{S}^n \to \mathbb{R}P^n$ given by sending a unit vector to its equivalence class. This is a double-covering map, and it induces a real projective structure on $\mathbf{S}^n$. The group $\mathsf{Aut}(\mathbf{S}^n)$ of projective automorphisms is isomorphic to the group $\mathsf{SL}_{\pm}(n+1, \mathbb{R})$ of linear maps of determinant equal to ± 1 with the quotient homomorphism $\mathsf{SL}_{\pm}(n+1, \mathbb{R}) \to \mathsf{PGL}(n+1, \mathbb{R})$ with the kernel $\pm\mathrm{I}$.

For an open domain $\mathcal{D}$ in $\mathbf{S}^n$ and a discrete group Γ in $\mathsf{Aut}(\mathbf{S}^n)$ acting on it properly discontinuously and freely, $\mathcal{D}/\Gamma$ has a real projective structure since $q_1|\mathcal{D}$ gives an immersion and the homomorphism

$$\Gamma \subset \mathsf{Aut}(\mathbf{S}^n) \to \mathsf{PGL}(n+1, \mathbb{R})$$

gives a holonomy homomorphism.

In this paper, we will study only $\mathbf{S}^3$. A homogeneous coordinate system of $\mathbf{S}^3$ is given by setting a point p of $\mathbf{S}^3$ be given coordinates $[t, x, y, z]$ where $\frac{(t,x,y,z)}{|(t,x,y,z)|}$ represents p as a unit vector in $\mathbf{S}^3$. Again

$$[t, x, y, z] \sim [t', z', y', z'] \text{ iff } (t, x, y, z) = \lambda(t', x', y', z') \text{ for } \lambda > 0.$$

There is the antipodal map

$$\mathscr{A} : \mathbf{S}^3 \to \mathbf{S}^3 \text{ given by } [\mathsf{v}] \to [-\mathsf{v}].$$

Given a subset or a point K of $\mathbf{S}^3$, we denote by K_- the set of antipodal points of points of K. Here, we denote by $\overline{pqp_-}$ the unique closed segment in $\mathbf{S}^3$ with endpoints p and p_- passing through q not equal to p or p_-. By $\overline{pq}$, we denote the unique closed segment in $\mathbf{S}^3$ with endpoints p and q provided $p \neq q$ and $p \neq q_-$.

A *subspace* of $\mathbf{S}^3$ is a subset defined by a system of linear equations in $\mathbb{R}^4$. A *line* is a 1-dimensional subspace. A singleton is not a subspace. A *projective geodesic* is an arc in a line. We will be using the standard Riemannian metric on $\mathbf{S}^3$ to be denoted by $\mathbf{d}_{\mathbf{S}^3}$. Notice that the geodesic structure of $\mathbf{S}^3$ is the same as the projective one. Thus, we will use the same term "geodesics" for both types of geodesics. Geodesics and totally geodesic subspaces are all subsets of lines or subspaces in $\mathbf{S}^3$. A projective automorphism sends these to themselves, while it does not preserve any metric. A pair of *antipodal points* on $\mathbf{S}^3$ is the equivalence class $[v]$ of a nonzero vector v and one $[-v]$ of $-v$. A segment connecting two antipodal points is precisely a segment of the $\mathbf{d}_{\mathbf{S}^3}$-length π.

The Lorentzian space $\mathsf{E}^{2,1}$ is the affine space $\mathbb{R}^3$ equipped with a nondegenerate bilinear form $\mathbf{B}$ of signature $1, 1, -1$. If we choose the origin in $\mathsf{E}^{2,1}$, we obtain the Lorentzian vector space $\mathsf{V}^{2,1}$.

One can think of $\mathbb{R}^3$ to be identical with $\mathsf{E}^{2,1}$ and the complement of a subspace of codimension-one in $\mathbb{RP}^3$, so called the complete affine subspace. Now $\mathbb{R}^3 = \mathsf{E}^{2,1}$ lifts a subspace of $\mathbf{S}^3$ double-covering $\mathbb{RP}^3$. The closure of the lifted $\mathsf{E}^{2,1}$ is a standard hemisphere $\mathscr{H}$, and $\mathsf{E}^{2,1}$ can be identified with the open hemisphere $\mathscr{H}^o$. The boundary $\mathbf{S}^2$ of $\mathsf{E}^{2,1}$ is the subspace of codimension-one identifiable with the projective 2-sphere $\mathbb{S}$ of directions in $\mathbb{R}^3$. (We identify $\mathsf{E}^{2,1}$ with the open affine space given by $t > 0$ in $\mathbf{S}^3$ from now on for convenience.)

The sphere $\mathbb{S}$ corresponds to the hyperplane given by $t = 0$ and the origin of $\mathsf{E}^{2,1}$ is given by $[1, 0, 0, 0]$, and by choosing O. For this system of coordinates, we have that the point $(x, y, z) \in \mathsf{E}^{2,1}$ is given coordinates $[1, x, y, z]$.

We will often identify $\mathbb{S}$ with the sphere of radius 1 in $\mathbb{R}^3$ and as a codimension-one totally geodesic subspace of $\mathbf{S}^3$. $\mathbb{S}$ also has a homogeneous coordinate system $[x, y, z]$ assigned to a vector p in $\mathbb{S}$ of unit length 1 if $\frac{(x,y,z)}{||(x,y,z)||} = p$. In the larger coordinate system $[x, y, z]$ is identical with $[0, x, y, z]$.

We will also be using a fixed Euclidean metric d_E on $\mathsf{E}^{2,1}$ compatible with the affine coordinate system. It is of great importance that projective geodesics, spherical geodesics, and Euclidean geodesics on $\mathsf{E}^{2,1}$ are the same ones up to parametrizations.

The group $\mathsf{Isom}^+(\mathsf{E}^{2,1})$ of orientation-preserving Lorentzian transformations of $\mathsf{E}^{2,1}$ is in the group $\mathsf{Aff}^+(\mathsf{E}^{2,1})$ of orientation-preserving affine transformations of $\mathsf{E}^{2,1}$. $\mathsf{Aff}^+(\mathsf{E}^{2,1})$ can be identified with the subgroup of $\mathsf{Aut}(\mathbf{S}^3)$ of orientation-preserving elements acting on $\mathbb{R}^3 = \mathsf{E}^{2,1}$, preserving $\mathbb{S}$, since any affine transformation of $\mathbb{R}^3$ extends to a projective automorphism of $\mathbb{R}\mathsf{P}^3$ and hence to an automorphism of $\mathbf{S}^3$ analytically. The elements of respective groups are called *projective automorphisms* of *Lorentzian* type and ones of *affine* type or just *Lorentzian isometries* or *affine transformations.* A *quasi-isometry* of $\mathbf{S}^3$ is a homeomorphism $f : \mathbf{S}^3 \to \mathbf{S}^3$ so that

$$C^{-1}\mathbf{d}_{\mathbf{S}^3}(x, y) \leq \mathbf{d}_{\mathbf{S}^3}(f(x), f(y)) \leq C\mathbf{d}_{\mathbf{S}^3}(x, y) \text{ for } x, y \in \mathbf{S}^3$$

where $C > 0$ is independent of x, y. Elements of $\mathsf{Aut}(\mathbf{S}^3)$ including ones extending the Lorentzian isometries are quasi-isometries of $\mathbf{S}^3$.

1.2. **Notation: Topology and geometry of projective $\mathbf{S}^3$.** We will denote by $\mathrm{bd}A$ or $\mathrm{bd}_X A$ the topological boundary of a subset A of some topological space X and by $\mathrm{int}A$ or $\mathrm{int}_X A$ the topological interior. If A is a topolgical manifold with boundary, we will denote the manifold boundary by ∂A and we will denote the manifold interior by A^o.

Given a pair of points or sets A and B, we define

$$\mathbf{d}_{\mathbf{S}^3}(A, B) = \inf\{\mathbf{d}_{\mathbf{S}^3}(x, y) | x \in A, y \in B\}.$$

An *ϵ-neighborhood* $N_\epsilon(A)$ for a number $\epsilon > 0$ of a subset A of $\mathbf{S}^3$ is the set of points of $\mathbf{d}_{\mathbf{S}^3}$-distances less than ϵ from some points of A.

We define the geometric Hausdorff distance $\mathbf{d}^H_{\mathbf{S}^3}(A, B)$ between two compact subsets A and B of $\mathbf{S}^3$ to be the infimum of

$$\{\epsilon \geq 0 | B \subset N_\epsilon(A) \text{ and } A \subset N_\epsilon(B)\}.$$

A sequence of compact sets $\{K_i\}$ in $\mathbf{S}^3$ *converges* or *converges geometrically* to a compact subset K if for every $\epsilon > 0$, there exists N so that for $i > N$, $\mathbf{d}^H_{\mathbf{S}^3}(K, K_i) < \epsilon$ or equivalently

$$K \subset N_\epsilon(K_i) \text{ and } K_i \subset N_\epsilon(K).$$

The space of compact subsets of a compact Hausdorff space with this metric is compact Hausdorff also and hence every sequence has a convergent subsequence.

1.3. **Real projective structures on surfaces.** Let ϑ be a projective automorphism $\mathsf{PGL}(3, \mathbb{R})$ represented by a diagonal matrix with distinct positive eigenvalues. Then it has three fixed points in $\mathbb{R}\mathsf{P}^2$: an attracting fixed point of the action of $\langle\vartheta\rangle$, a repelling fixed point, and a saddle-type fixed point. Three lines passing through pairs of them are ϑ-invariant, as are four open triangles in the complement of the union of these. The components of the lines with fixed points removed are so-called open sides. The union of two open sides of one of the open triangles ending at an attracting fixed point or a repelling fixed point simultaneously is acted properly and freely upon by $\langle\vartheta\rangle$. The quotient space is diffeomorphic to a compact annulus. The real projective surface projectively diffeomorphic to the quotient space is said to be an *elementary annulus*.

Take two adjacent ϑ-invariant triangles among the four triangles with three open sides of them all ending in an attracting fixed point or a repelling fixed point. Then the quotient of the union by $\langle\vartheta\rangle$ is diffeomorphic to an annulus. A projectively diffeomorphic real projective surface is said to be a π-*annulus*. (See [10] and [11] for more details.)

Let $\tilde{S}$ be the universal cover of a real projective surface S with a developing map $\mathsf{dev} : \tilde{S} \to \mathbb{R}\mathsf{P}^2$ and a holonomy homomorphism h. For a circle $\mathbf{S}^1$, a *closed geodesic* $c : \mathbf{S}^1 \to S$ is a closed curve where given a lift $\tilde{c} : \mathbb{R} \to \tilde{S}$ of c, $\mathsf{dev} \circ \tilde{c}$ is a straight arc in $\mathbb{R}\mathsf{P}^2$. A closed geodesic is *principal* if for a lift $\tilde{c}$ to the universal cover $\tilde{S}$, $\mathsf{dev} \circ \tilde{c}$ is an imbedding to a straight arc connecting an attracting and a repelling fixed point of $h(\gamma)$ for a deck transformation γ of $\tilde{S}$ satisfying $\tilde{c}(t + 2\pi) = \gamma \circ \tilde{c}(t)$. (Here γ is said to be the *corresponding* deck transformation of $\tilde{c}$ and c and it exists uniquely.)

A *properly convex* domain in $\mathbb{R}\mathsf{P}^2$ is a bounded convex domain of an affine subspace in $\mathbb{R}\mathsf{P}^2$. A real projective surface is *properly convex* if it is a quotient of a properly convex domain in $\mathbb{R}\mathsf{P}^2$ by a properly discontinuous and free action of a subgroup of $\mathsf{PGL}(3, \mathbb{R})$.

A disjoint collection of simple closed geodesics $c_1, \ldots, c_m$ *decomposes* a real projective surface S into subsurfaces $S_1, .., S_n$ if each S_i is the closure of a component of $S - \bigcup_{i=1,..,m} c_i$ where we do not allow a curve c_i to have two one-sided neighborhoods in only one S_i for some i.

In [11], we proved:

Theorem 3 (Choi). *Let Σ be a closed orientable real projective surface with principal geodesic or empty boundary and $\chi(\Sigma) < 0$. Then Σ has a collection of disjoint simple closed principal geodesics decomposing Σ into properly convex real projective surfaces with principal geodesic boundary and of negative Euler characteristic and/or π-annuli with principal geodesic boundary.*

1.4. **Joins.** Suppose that two compact subsets A and B of $\mathbf{S}^3$ have no pairs of points $x \in A, y \in B$ so that x and y are antipodal. Then the *join* $J(A, B)$ is the union of all segments of $\mathbf{d}_{\mathbf{S}^3}$-lengths $< \pi$ with single endpoints in A and the other ones in B. In a set theoretic terms,

$$(1)\quad J(A,B) = q_1(\{c_1\vec{u}_1 + c_2\vec{u}_2 | c_1, c_2 \geq 0, \vec{u}_1 \in q_1^{-1}(A), \vec{u}_2 \in q_1^{-1}(B)\}) \subset \mathbb{R}^4.$$

Normally, $A \cup B$ is required not to have antipodal points here. However, we later need exceptional cases when one of them A is a great circle and B is a point and every segment from B to A is of $\mathbf{d}_{\mathbf{S}^3}$-length $< \pi - \epsilon$ for a fixed $\epsilon > 0$. In this case, $J(A, B)$ is a *hemisphere*, a totally geodesic disk with geodesic boundary in $\mathbf{S}^3$, a convex set.

If A and B are convex and $A \cup B$ has no pair of antipodal points, then $J(A, B)$ is also properly convex. The proof can be seen by reverting back to $\mathbb{R}^4$ by equation 1.

Lemma 4. *Let A_i and B_i be a sequence of properly convex compact subsets of $\mathbf{S}^3$ geometrically converging to properly convex compact sets A and B respectively. Suppose that $A \cup B$ has no antipodal pair of points. Then the sequence $\{J_i\}$ of the join J_i of A_i and B_i for each i converges to the join J of A and B.*

Proof. Clearly, A_i and B_i have no antipodal pair of points for sufficiently large i. Now this is straightforward. □ □

Lemma 5. *Let $\{h_i\}_{i \in \mathbb{Z}_+}$ be a sequence of hemispheres in $\mathbf{S}^3$ with ∂h_i converging to a great circle $\mathbf{S}^1_0$ and suppose that a sequence of points $\{p_i | p_i \in h_i^o\}_{i \in \mathbb{Z}_+}$ converges to a point p_∞ not in $\mathbf{S}^1_0$. Then h_i converges to the hemisphere h_∞ with boundary $\mathbf{S}^1_0$ containing p_∞.*

Proof. The hemisphere h_i is a join of p_i with ∂h_i and is a union of segments from p_i to points on ∂h_i. Since p_∞ is not antipodal to points of $\mathbf{S}^1_0$, the result follows. □ □

1.5. **Oriented Lorentzian vector spaces.** Let $(\mathsf{V}^{2,1}, \cdot, \mathsf{Det})$ denote a three-dimensional *oriented Lorentzian* $\mathbb{R}$-vector space. That is, $\mathsf{V}^{2,1} \cong \mathbb{R}^3$ and is given a symmetric bilinear form

$$\mathbf{B} : \mathsf{V}^{2,1} \times \mathsf{V}^{2,1} \longrightarrow \mathbb{R}$$

defined by $(\mathsf{v},\mathsf{u}) \mapsto \mathsf{v}\cdot\mathsf{u}$ of index 1 and a nondegenerate alternating trilinear form

$$\mathsf{V}^{2,1} \times \mathsf{V}^{2,1} \times \mathsf{V}^{2,1} \longrightarrow \mathbb{R}$$

defined by $(\mathsf{v},\mathsf{u},\mathsf{w}) \mapsto \mathsf{Det}(\mathsf{v},\mathsf{u},\mathsf{w})$. Its automorphism group is the special orthogonal group $\mathsf{SO}(2,1)$.

The Lorentzian structure divides $\mathbb{S}$ into three open domains $\mathbb{S}_+, \mathbb{S}_0, \mathbb{S}_-$ separated by two conics $\mathrm{bd}\mathbb{S}_+$ and $\mathrm{bd}\mathbb{S}_-$. Let $\mathbb{S}_+ \subset \mathbb{S}$ denote the set of future time-like directions; its boundary $\mathrm{bd}\mathbb{S}_+$ consists of future null directions. Similarly, let $\mathbb{S}_- \subset \mathbb{S}$ denote the set of past time-like directions with boundary $\mathrm{bd}\mathbb{S}_-$ consisting of past null directions. Both $\mathbb{S}_+$ and $\mathbb{S}_-$ are cells, interchanged by $\mathscr{A}$. The set $\mathbb{S}_0$ of space-like directions forms an $\mathscr{A}$-invariant annulus bounded by the disjoint union $\mathrm{bd}\mathbb{S}_+ \coprod \mathrm{bd}\mathbb{S}_-$.

Recall that $\mathbb{S}_+$ is the Beltrami-Klein model of the hyperbolic plane where $\mathsf{SO}(2,1)$ acts as the orientation-preserving isometry group. Here the metric geodesics are precisely the projective geodesics and vice versa. (We will use the same term "geodesics" for both types of geodesics for ones in $\mathbb{S}_+$ and later $\mathbb{S}_-$.) The geodesics in $\mathbb{S}_+$ are straight arcs and $\mathrm{bd}\mathbb{S}_+$ forms the ideal boundary of $\mathbb{S}_+$. For a finitely generated discrete subgroup Γ in $\mathsf{SO}(2,1)$, $\mathbb{S}_+/\Gamma$ has a complete hyperbolic structure as well as a real projective structure with the compatible geodesic structure.

Alternatively, we can use $\mathbb{S}_-$ as the model and $\mathbb{S}_-/\Gamma$ has a complete hyperbolic structure as well as a real projective structure. Since $\mathbb{S}_+$ and $\mathbb{S}_-$ are properly convex, these are examples of properly convex real projective surfaces.

1.6. **Null half-planes.** Let $\mathcal{N}$ denote the *nullcone* in $\mathsf{V}^{2,1}$, that is, the collection of all vectors $\mathsf{v} \in \mathsf{V}^{2,1}$ with $\mathsf{v}\cdot\mathsf{v} = 0$. Its projectivization $\mathsf{P}(\mathcal{N} - \{O\})$ consists of all *null lines* in $\mathsf{V}^{2,1}$. If $\mathsf{v} \in \mathcal{N} - \{O\}$, then its orthogonal complement $\mathsf{v}^\perp$ is a *null plane* which contains the line $\mathbb{R}\mathsf{v}$. The line $\mathbb{R}\mathsf{v}$ separates $\mathsf{v}^\perp$ into two half-planes.

The orientation on $\mathsf{V}^{2,1}$ determines an $\mathsf{SO}(2,1)$-invariant way to uniquely associate a component of $\mathsf{v}^\perp - \mathbb{R}\mathsf{v}$ to $\mathbb{R}\mathsf{v}$ as follows. Since $\mathsf{v} \in \mathcal{N}$, its direction lies in either $\mathrm{bd}\mathbb{S}_+$ or $\mathrm{bd}\mathbb{S}_-$. Choose an arbitrary element u of $\mathbb{S}_+$ or $\mathbb{S}_-$ respectively, so that the directions of v and u both lie in $\mathrm{Cl}(\mathbb{S}_+)$ or $\mathrm{Cl}(\mathbb{S}_-)$ respectively. (For example $\mathsf{u} = (0,0,\pm1)$ would be sufficient.) Define the *null half-plane* $\mathscr{W}(\mathsf{v})$ (or the *wing*) associated to v as:

$$\mathscr{W}(\mathsf{v}) := \{\mathsf{w} \in \mathsf{v}^\perp \mid \mathsf{Det}(\mathsf{v},\mathsf{w},\mathsf{u}) > 0\}.$$

Since the Lorentzian product $\mathsf{u} \otimes \mathsf{v}$ is in $\mathscr{W}(\mathsf{v})$, the directions in the wing satisfy the right-hand rule. Since $\mathscr{W}(\mathsf{v}) = \mathscr{W}\big(\mathscr{A}(\mathsf{v})\big)$, the null half-plane $\mathscr{W}(\mathsf{v})$ depends only on $\mathbb{R}\mathsf{v}$. The corresponding set of directions is the open arc

$$\varepsilon(\mathsf{v}) = \varepsilon([\mathsf{v}]) := [\mathscr{W}(\mathsf{v})]$$

in $\mathbb{S}_0$ joining $[\mathsf{v}]$ to $\mathscr{A}([\mathsf{v}]) = [-\mathsf{v}]$. Since $\mathsf{v}^\perp$ is tangent to $\mathcal{N}$, the arc $\varepsilon(\mathsf{v})$ is tangent to $\mathrm{bd}\mathbb{S}_+$. The orientation of $\mathrm{bd}\mathbb{S}_+$ as the boundary orientation of $\mathbb{S}_+$ agrees with the orientation of $\varepsilon(\mathsf{v})$ away from $[\mathsf{v}]$.

The corresponding map $[\mathsf{v}] \longmapsto \varepsilon(\mathsf{v})$ is an $\mathsf{SO}(2,1)$-equivariant map

$$\mathrm{bd}\mathbb{S}_+ \to \mathcal{S}$$

where $\mathcal{S}$ denotes the set of half-arcs of form $\varepsilon(\mathsf{v})$ for $\mathsf{v} \in \mathrm{bd}\mathbb{S}_+$. The arcs $\varepsilon([\mathsf{v}])$ for $\mathsf{v} \in \mathrm{bd}\mathbb{S}_+$ foliate $\mathbb{S}_0$ as we can see from an $\mathbf{S}^1$-action fixing $[\pm 1, 0, 0]$ where $\mathbf{S}^1$ is a subgroup of $\mathsf{SO}(2,1)$. Let us call the foliation $\mathcal{F}$.

Hence $\mathbb{S}_0$ has a $\mathsf{SO}(2,1)$-equivariant quotient map

$$\Pi : \mathbb{S}_0 \to \mathsf{P}(\mathcal{N} - \{O\}) \cong \mathbf{S}^1$$

and $\varepsilon([\mathsf{v}]) = \Pi^{-1}([\mathsf{v}])$ for each $\mathsf{v} \in \mathcal{N} - \{O\}$.

Consider a future-pointing null vector of Euclidean length $\sqrt{2}$:

$$\mathsf{n}_\theta := \begin{bmatrix} \cos(\theta) \\ \sin(\theta) \\ 1 \end{bmatrix} \text{ for } \theta \in [0, 2\pi)$$

Then $\mathscr{W}(\mathsf{n}_\theta)$ consists of all

$$\mathsf{p}_\theta(t,s) := t\mathsf{n}_\theta + s \begin{bmatrix} -\sin(\theta) \\ \cos(\theta) \\ 0 \end{bmatrix}$$

where $t \in \mathbb{R}$ and $s > 0$. The arc $c_\theta := \varepsilon(\mathsf{n}_\theta)$ on the sphere of directions $\mathbb{S}$ is parametrized by as unit vectors

$$\frac{1}{\sqrt{2}}\mathsf{p}_\theta(t, \sqrt{1-t^2})$$

as $-1 < t < 1$. (As an element of $\mathbf{S}^3$, its homogeneous coordinates equal

$$[0, \frac{1}{\sqrt{2}}\mathsf{p}_\theta(t, \sqrt{1-t^2})]$$

where $\frac{1}{\sqrt{2}}\mathsf{p}_\theta(t, \sqrt{1-t^2})$ is used as the last three coordinates.)

The other component of $\mathsf{v}^\perp - \mathbb{R}\mathsf{v}$ could also be used as a wing $\mathscr{W}(\mathsf{v})$, but for the opposite orientation on $\mathsf{V}^{2,1}$. Alternatively, these *negatively oriented null half-planes* are the images of the positively oriented null half-planes under the antipodal map $\mathscr{A}$ (which reverses orientation in dimension three). This phenomenon appears in the theory of crooked planes, where one must choose a class of orientations for crooked planes, as well as the Margulis invariant, where the choice of orientation plays a crucial role.

Since we have no need to consider negatively oriented null half-planes, we henceforth restrict our attention to positively oriented null half-planes in this paper.



Figure 1. The tangent geodesics to disks $\mathbb{S}_+$ and $\mathbb{S}_-$ in the unit sphere $\mathbb{S}$ imbedded in $\mathbb{R}^3$.



Figure 2. The tangent geodesics to disks $\mathbb{S}_+$ and $\mathbb{S}_-$ in the stereographically projected $\mathbb{S}$ from $(0, 0, -1)$. Radial arcs are geodesics. The inner circle represents the boundary of $\mathbb{S}_+$. The arcs of form $\varepsilon(x)$ for $x \in \mathrm{bd}\mathbb{S}_+$ are leaves of the foliation $\mathcal{F}$ on $\mathbb{S}_0$.

2. The real projective surface that is a quotient of a domain in $\mathbb{S}$.

Goldman constructed a real projective surface Σ with a free holonomy group in [24]. The holonomy group is in the image $\mathsf{PSO}(2, 1)$ of $\mathsf{SO}(2, 1)$ in $\mathsf{PGL}(3, \mathbb{R})$. We will show that a domain in $\mathbb{S}$ regularly covers Σ.

2.1. **The construction of the Γ-invariant domain.**

As above, let Γ be an orientation-preserving finitely generated Lorentzian isometry group without any parabolic elements acting freely and properly discontinuously on $\mathbb{S}_+$ (or equivalent on $\mathbb{S}_-$) isomorphic to a free group of finite rank ≥ 2. Then $\mathbb{S}_+/\Gamma$ is a complete genus $\tilde{\mathsf{g}}$ hyperbolic surface with b ideal boundary components where $\mathsf{b} \geq 1$ and $\tilde{\mathsf{g}}$ is just some integer ≥ 0. In other words, $\mathbb{S}_+/\Gamma$ can be compactified by adding boundary components to a compact surface Σ'_+ with b boundary components. (Σ'_+ itself is a real projective surface with boundary.) The universal cover $\tilde{\Sigma}'_+$ of Σ'_+ is identified with a domain $\mathbb{S}_+$ union a collection of open arcs in $\mathrm{bd}\mathbb{S}_+$. (At the moment, we do not require that Γ acts properly discontinuously or freely on $\mathsf{E}^{2,1}$.)

Let Λ denote the limit set of an orbit $\Gamma(x)$ for $x \in \mathbb{S}_+$, which is independent of the choice of x, as is well-known from the classical Fuchsian group theory. We know from the Fuchsian group theory that Λ is a Cantor subset of $\mathrm{bd}\mathbb{S}_+$ and of Lebesgue measure zero and the set of fixed points of nonidentity elements of Γ is dense in Λ. We can identify $\tilde{\Sigma}'_+$ as $\mathrm{Cl}(\mathbb{S}_+) - \Lambda$. (See Chapter 8 of Beardon [3] or Chapter 2 of Marden [32] for the classical Fuchsian group theory used here.)

We will be using the same symbols for Γ its extension to $\mathbf{S}^3$ and its element g and its extension to $\mathbf{S}^3$. We identify $\mathbb{S}$ with the boundary of $\mathscr{H}$, called a *boundary sphere.* Γ acts on the standard circles $\mathrm{bd}\mathbb{S}_\pm$ and the interiors $\mathbb{S}_\pm$ in $\mathbb{S}$. Then each nonidentity element g of Γ has an attracting fixed point a in $\mathrm{bd}\mathbb{S}_+$, a repelling fixed point r in $\mathrm{bd}\mathbb{S}_+$, and a saddle-type fixed point s in $\mathbb{S}_0$. Of course, their antipodes in $\mathbb{S}$ are also an attracting fixed point a_- in $\mathrm{bd}\mathbb{S}_-$ and a repelling fixed point r_- in $\mathrm{bd}\mathbb{S}_-$ and a saddle-type fixed point s_- in $\mathbb{S}_0$ respectively.

Since g acts on $\mathrm{bd}\mathbb{S}_+$ and fixes a and r, we obtain that g acts on the two 1-dimensional subspaces tangent to $\mathrm{bd}\mathbb{S}_+$ at a and r. Hence, $\{s, s_-\}$ is their intersection. Therefore, we deduce that

$$\{s, -s\} \subset \varepsilon(a) \cup \varepsilon(r) \tag{2}$$

as $\varepsilon(a)$ and $\varepsilon(r)$ are disjoint halves of these 1-dimensional subspaces.

Let $\mathcal{J}$ be the set indexing the boundary components of $\tilde{\Sigma}'$. We denote the boundary components of $\tilde{\Sigma}'$ by $\mathbf{b}_i, i \in \mathcal{J}$. We obtain a convex domain Ω_+ closed in $\mathbb{S}_+$ bounded by a union of straight segments $\mathbf{l}_i$ for $i \in \mathcal{J}$ with same endpoints as $\mathbf{b}_i$. That is, $\mathbf{l}_i$ are geodesics in $\mathbb{S}_+$ connecting ideal endpoints of $\mathbf{b}_i$. Here, $\mathbf{l}_i$ is $\langle \mathbf{g}_i \rangle$-invariant for a unique primitive element of $\mathbf{g}_i \in \Gamma$ where $\mathbf{g}_i$ acts along the orientation of $\mathbf{l}_i$. Since Γ acts a geometrically finite Fuchsian group on $\mathbb{S}_+$ without parabolics, the classical Fuchsian group theory tells us that Ω_+/Γ is a compact hyperbolic surface with geodesic boundary that is the convex hull of $\mathbb{S}_+/\Gamma$ and homeomorphic to Σ'_+.

We now denote by $\tilde{\Sigma}'_-$ the domain of antipodal points of $\tilde{\Sigma}'$ in $\mathbb{S}_-$; that is, $\tilde{\Sigma}'_- = \mathscr{A}(\tilde{\Sigma}')$. We note that Γ acts properly discontinuously and freely on $\tilde{\Sigma}'_-$ by antipodality, and $\tilde{\Sigma}'_-/\Gamma$ is a compact real projective surface with geodesic boundary diffeomorphic to Σ'_+. Denote it by Σ'_-. Also, let Ω_- be the set of antipodal points to Ω_+. Then Γ acts properly discontinuously and freely on Ω_+ and Ω_-. Denote the respective compact quotient surfaces with geodesic boundary by Σ''_+ and Σ''_-.

For each point x of $\mathrm{bd}\mathbb{S}_+$, we recall that $\varepsilon(x)$ is an open segment of length π with end points x and $-x$ tangent to $\mathrm{bd}\mathbb{S}_+$. The open arcs $\varepsilon(x)$ for $x \in \mathrm{bd}\mathbb{S}_+$ give leaves of the foliation $\mathcal{F}$ in $\mathbb{S}_0$. Let us give an induced orientation on Σ'_+ and hence an induced orientation on each $\mathbf{l}_i$ for $i \in \mathcal{J}$. Let p_i and q_i denote the forward and backward end points of $\mathbf{l}_i$. We draw a segment $s_i = \varepsilon(p_i)$. We also draw a segment $t_i = \varepsilon(q_i)$. Then $\mathbf{l}_i, s_i, t_i, \mathbf{l}_{i,-}$ bound an open disk invariant under $\langle \mathbf{g}_i \rangle$, which we call a *strip*. We denote by $\mathcal{R}_i$ the open strip union with $\mathbf{l}_i$ and $\mathbf{l}_{i,-}$. (See Figures 1 and 2 for the pictures of these arcs tangent to $\mathbb{S}_+$ and $\mathbb{S}_-$.)

Using $\mathcal{F}$, we obtain:

Proposition 6. *The strips $\mathcal{R}_i$ and $\mathcal{R}_j$ are disjoint for $i \neq j$ where $i, j \in \mathcal{J}$.*

Proof. Since s_i, t_i, s_j, t_j are all leaves of $\mathcal{F}$ corresponding to distinct points of $\mathrm{bd}\mathbb{S}_+$, we obtain the disjointness. (See Figure 3.) We have that s_i does not intersect t_i, s_j, t_j for $i \neq j$, $i, j \in \mathcal{J}$, and t_i does not intersect s_i, s_j, t_j for $i \neq j$. The elements $\mathbf{l}_i$ and $\mathbf{l}_{i,-}$ for $i \in \mathcal{J}$ are mutually disjoint. Thus the boundary $\mathrm{bd}\mathcal{R}_i$ and $\mathrm{bd}\mathcal{R}_j$ for $i \neq j$ are disjoint. Hence, the conclusion follows. □ □

Goldman in his thesis classified compact annuli with geodesic boundary and holonomy matrices that are diagonalizable with distinct positive eigenvalues [23] following Nagano-Yagi [35]. (See [11] also.)



FIGURE 3. The quotient spaces of the two "lune" regions are π-annuli.

Proposition 7. *Given* $\mathcal{R}_i$ *for* $i \in \mathcal{J}$ *and* $\gamma \in \Gamma$, *we either have*

$$\gamma(\mathcal{R}_i) = \mathcal{R}_i \text{ or } \gamma(\mathcal{R}_i) \cap \mathcal{R}_i = \emptyset.$$

In the former case, $\gamma = \mathbf{g}_i^n$, $n \in \mathbb{Z}$ *for the deck transformation* $\mathbf{g}_i$ *corresponding to* $\mathbf{l}_i$. *Furthermore,* $(\mathcal{R}_i \cup \mathbf{l}_i \cup \mathbf{l}_{i,-})/\langle \mathbf{g}_i \rangle$ *is a compact annulus with geodesic boundary that decomposes into two* π*-annuli or four elementary annuli along some collection of simple closed geodesics.*

Proof. If we have $\mathcal{R}_i \cap \gamma(\mathcal{R}_i) \neq \emptyset$, then we obtain $\mathbf{l}_i \cap \gamma(\mathbf{l}_i) \neq \emptyset$. Since $\mathbf{l}_i$ goes to boundary components of Ω_+/Γ, it follows that $\mathbf{l}_i = \gamma(\mathbf{l}_i)$ and $\mathcal{R}_i = \gamma(\mathcal{R}_i)$ and $\gamma = \mathbf{g}_i^n$ for $n \in \mathbb{Z}$.

There is a subspace l of dimension 1, a geodesic circle, containing $\mathbf{l}_i$ and $\mathbf{l}_{i,-}$ in $\mathbf{S}^2$. l and t_i bound a closed disk D_1 and if we remove t_i and the fixed points at the ends of $\mathbf{l}_i$, we see that this region covers a π-annulus with principal boundary components. The similar statement is true for the disk bounded by l and s_i. Therefore, $A_i := (\mathcal{R}_i \cup \mathbf{l}_i \cup \mathbf{l}_{i,-})/\langle \mathbf{g}_i \rangle$ is a union of two compact π-annuli. In other words, $l - \mathrm{Cl}(\mathbf{l}_i) - \mathrm{Cl}(\mathbf{l}_{i,-})$ maps to a simple closed principal geodesic decomposing A_i into two π-annuli with principal boundary. (See Figure 3.) Also, each π-annulus decomposes into two elementary annuli. (See Section 1.3.) □ □

We say that for $i, j \in \mathcal{J}$, the annulus $\mathcal{R}_i/\langle \mathbf{g}_i \rangle$ is *equivalent* to $\mathcal{R}_j/\langle \mathbf{g}_j \rangle$ if $\mathcal{R}_j = g(\mathcal{R}_i)$ and $g\mathbf{g}_i g^{-1} = \mathbf{g}_j$ for $g \in \Gamma$. Thus, in fact, there are only b equivalence classes of annuli of above form.

We define $\mathcal{A}_i = \mathcal{R}_i \cap \mathbb{S}_0$ for $i \in \mathcal{J}$, which equals $\bigcup_{x \in \mathbf{b}_i} \varepsilon(x)$. We note that $\mathcal{A}_i \subset \mathcal{R}_i$ for each $i \in \mathcal{J}$. We finally define

$$\begin{aligned}
\tilde{\Sigma} &= \tilde{\Sigma}'_+ \cup \coprod_{i \in \mathcal{J}} \mathcal{R}_i \cup \tilde{\Sigma}'_- \\
&= \tilde{\Sigma}'_+ \cup \coprod_{i \in \mathcal{J}} \mathcal{A}_i \cup \tilde{\Sigma}'_- \\
&= \Omega_+ \cup \coprod_{i \in \mathcal{J}} \mathcal{R}_i \cup \Omega_- \qquad (3) \\
&= \mathbb{S} - \bigcup_{x \in \Lambda} \mathrm{Cl}(\varepsilon(x)). \qquad (4)
\end{aligned}$$

an open domain in $\mathbb{S}$. Since the collection whose elements are of form $\mathcal{R}_i$ mapped to itself by Γ, we showed that Γ acts on this open domain.

2.2. **The real projective surface.** First, $\tilde{\Sigma}$ does not contain any fixed point: Suppose that $g \in \Gamma$ acts on $\mathbf{l}_i$ for some $i \in \mathcal{J}$. Then g acts on $\mathcal{R}_i$ and the four attracting and repelling fixed points are the vertices of $\mathcal{R}_i$ and two saddle-type fixed points are in s_i or t_i. Hence, they are outside $\tilde{\Sigma}$.

Suppose that $g \in \Gamma - \{\mathrm{I}\}$ does not act on any of the boundary components. Then it has four attracting and repelling fixed points $a, r \in \mathrm{bd}\mathbb{S}_+$ and $a_-, r_- \in \mathrm{bd}\mathbb{S}_-$. By equation 2, saddle-type fixed points s and s_- are either on a great segment $\varepsilon(a)$ or the other one $\varepsilon(r)$ tangent to $\mathrm{bd}\mathbb{S}_+$ at a



Figure 4. A figure representing an actual free group action with some of the arcs s_i and t_i drawn in point-plots on stereographically projected sphere as in Figure 2.

and r respectively. Since we have $a, r \in \Lambda$, the fixed points are outside $\tilde{\Sigma}$ by equation 4.

For a subgroup $G \subset \mathsf{SO}(2,1)$ whose elements are orientation-preserving on $\mathbb{S}_+$, and any open domain $\mathcal{D}$ where G acts freely and properly-discontinuously on, $\mathcal{D}/G$ has an induced orientation from $\mathbb{S}_+$.

Theorem 8. *Let Γ be an orientation-preserving Lorentzian isometry group without any parabolic elements acting freely and properly discontinuously on $\mathbb{S}_+$ (or equivalent on $\mathbb{S}_-$) isomorphic to a free group of finite rank ≥ 2. Let b be the number of ideal boundary components of $\mathbb{S}_+/\Gamma$. Then*

- *there exists an open domain $\mathcal{D}$ in $\mathbb{S}$ where Γ acts properly discontinuously and freely, and $\mathcal{D}/\Gamma$ is homeomorphic to a closed surface of genus g, $\mathsf{g} \geq 2$.*

- *As a real projective surface $\mathcal{D}/\Gamma$ decomposes along simple closed principal geodesics into a union of b annuli $A_1, \dots, A_{\mathsf{b}}$ and two convex real projective surfaces Σ''_+ and Σ''_-. Each annulus A_i, $i = 1, \dots, \mathsf{b}$ decomposes into two π-annuli.*
- *Γ also acts properly discontinuously and freely on $\mathcal{D}_-$ and $\mathcal{D}_-/\Gamma$ is a closed real projective surface diffeomorphic to $\mathcal{D}/\Gamma$.*
- *$\mathcal{D}_-$ also decomposes into Σ''_+ and Σ''_- and b annuli A'_i.*
- *The antipodal map $\mathscr{A} : \mathbb{S} \to \mathbb{S}$ induces a real projective diffeomorphism between $\mathcal{D}/\Gamma$ and $\mathcal{D}_-/\Gamma$.*

Proof. We begin by letting $\mathcal{D} = \tilde{\Sigma}$ obtained as above. We showed that Γ has no fixed point on $\mathcal{D}$. Let $\mathrm{Cl}_{\mathcal{D}}(\mathcal{A}_i)$ denote the closure of $\mathcal{A}_i$ in $\mathcal{D}$ for each $i \in \mathcal{J}$.

Now Γ acts properly discontinuously on $\tilde{\Sigma}'_+$ and $\tilde{\Sigma}'_-$ and each $g \in \Gamma$ sends $\mathrm{Cl}_{\mathcal{D}}(\mathcal{A}_i)$ to $\mathrm{Cl}_{\mathcal{D}}(\mathcal{A}_j)$ for $j \neq i$ or g is in the infinite cyclic subgroup of Γ acting properly discontinuously on $\mathrm{Cl}_{\mathcal{D}}(\mathcal{A}_i)$ if $g(\mathrm{Cl}_{\mathcal{D}}(\mathcal{A}_i)) = \mathrm{Cl}_{\mathcal{D}}(\mathcal{A}_i)$. Hence, Γ acts properly discontinuously on $\bigcup_{i \in \mathcal{J}} \mathrm{Cl}_{\mathcal{D}}(\mathcal{A}_i)$.

For any compact set K in $\tilde{\Sigma}$, we consider the compact sets $K_1 := K \cap \tilde{\Sigma}'_+$, $K_{-1} := K \cap \tilde{\Sigma}'_-$, and $K_0 := K \cap \bigcup_{i \in \mathcal{J}} \mathrm{Cl}_{\mathcal{D}}(\mathcal{A}_i)$ since these are closed. We have $K = K_+ \cup K_- \cup K_0$. Note that K_0 meets only finitely many $\mathrm{Cl}_{\mathcal{D}}(\mathcal{A}_i)$.

If $g(K) \cap K \neq \emptyset$, we have $g(K_i) \cap K_i \neq \emptyset$ for some $i = -1, 0, 1$. Since Γ acts properly discontinuously on $\tilde{\Sigma}'_+$ and $\tilde{\Sigma}'_-$ and $\bigcup_{i \in \mathcal{J}} \mathrm{Cl}_{\mathcal{D}}(\mathcal{A}_i)$ respectively, there exist only finitely many element g of Γ so that $g(K_i) \cap K_i \neq \emptyset$ for each i, $i = -1, 0, 1$. We showed that Γ acts properly discontinuously and freely on $\mathcal{D}$.

For the second item, we see that $\mathcal{D}/\Gamma$ decomposes along the images of the union of $\mathbf{l}_i, \mathbf{l}_{i,-}, i \in \mathcal{J}$ to Σ''_+ and Σ''_- and annuli bounded by images of the union of $\mathbf{l}_i$ and $\mathbf{l}_{i,-}$, $i \in \mathcal{J}$. We see from Proposition 7 that these annuli are obtained from two π-annuli.

The third item follows since the antipodal map $\mathscr{A}$ conjugates Γ to itself and $\mathscr{A}(\mathcal{D}) = \mathcal{D}_-$. Let κ denote the induced diffeomorphism $\Sigma \to \mathcal{D}_-/\Gamma$ where κ sends Σ''_- to Σ''_+ and vice versa. κ sends annuli in Σ to ones in $\mathcal{D}_-/\Gamma$ but they do not share subdomains in the universal cover. (Actually $\mathcal{R}_i$ goes to its antipodal image which is distinct from $\mathcal{R}_i$ for each $i \in \mathcal{J}$.)

The rest follows from this observation. □ □

3. The work of Goldman, Labourie and Margulis

In this section, we will briefly sketch the relevant results of Goldman, Labourie, and Margulis in [28] and [27].

Let Γ be a nonamenable orientation-preserving Lorentzian isometry group without any parabolic elements acting freely and properly discontinuously on $\mathsf{E}^{2,1}$ and hence on $\mathbb{S}_+$ (or equivalent on $\mathbb{S}_-$) isomorphic to a free group of finite rank ≥ 2. Then $\mathbb{S}_+/\Gamma$ is a complete genus $\tilde{\mathsf{g}}$ hyperbolic surface with b ideal boundary components. For convenience, denote by Σ_+ the interior of the surface Σ'_+ from now on. That is, $\Sigma_+ = \mathbb{S}_+/\Gamma$.

3.1. **Margulis invariants.** Given an element $g \in \Gamma$, let us denote by $\mathsf{v}_+(g)$, $\mathsf{v}_0(g)$, and $\mathsf{v}_-(g)$ the eigenvectors of the linear part $\mathcal{L}(g)$ of g corresponding to eigenvalues > 1, $= 1$, and < 1 respectively. $\mathsf{v}_+(g)$ and $\mathsf{v}_-(g)$ are null vectors and $\mathsf{v}_0(g)$ is space-like and of unit norm. We choose so that $\{\mathsf{v}_+(g), \mathsf{v}_0(g), \mathsf{v}_-(g)\}$ are in the standard orientation of $\mathbb{R}^3$; or $\mathsf{v}_-(g) \otimes \mathsf{v}_+(g) = \mathsf{v}_0(g)$. We recall the Margulis invariant $\mu : \Gamma \rightarrow \mathbb{R}$

$$\mu(g) := \mathbf{B}(gx - x, \mathsf{v}_0(g)) \text{ for } g \in \Gamma, x \in \mathsf{E}^{2,1},$$

which is independent of the choice of x in $\mathsf{E}^{2,1}$. (See [18] for details.)

If Γ acts freely on $\mathsf{E}^{2,1}$, then Margulis invariants of nonidentity elements are all positive or all negative by the Opposite sign-lemma of Margulis (See [28].) We can change the all signs of the Margulis invariants by changing the orientation of $\mathsf{E}^{2,1}$. From now on, we assume that the Margulis invariants of nonidentity elements are all positive.

Let $\mathbb{US}_+$ be the unit tangent vector bundle over $\mathbb{S}_+$, and let $\mathbb{U}(\mathbb{S}_+/\Gamma) = \mathbb{U}\Sigma_+$ be the unit tangent vector bundle over $\mathbb{S}_+/\Gamma = \Sigma_+$. A *geodesic current* is a Borel probability measure on $\mathbb{U}(\mathbb{S}_+/\Gamma)$ which is supported on a union of weakly recurrent geodesics. We denote by $\mathcal{C}(\mathbb{S}_+/\Gamma)$ the space of geodesic currents supported on unions of closed geodesics which is a dense subset. In the second main theorem of [28], they extend the Margulis invariants to the geodesic current space on the hyperbolic space $\mathbb{S}_+/\Gamma$, i.e., diffused Margulis invariants for currents.

The Margulis invariants of elements of $\mathcal{C}(\mathbb{S}_+/\Gamma)$ are positive by the second main theorem in [28], and the subspace of currents supported closed geodesics with total measure 1 is precompact in $\mathcal{C}(\mathbb{S}_+/\Gamma)$. It follows that there exists a global $C > 0$ independent of $g \in \Gamma$ so that

$$\frac{\mu(g)}{l_{\mathbb{S}_+}(g)} \geq C \text{ for } g \in \Gamma - \{\mathrm{I}\} \tag{5}$$

where $l_{\mathbb{S}_+}(g)$ denotes the minimum of the hyperbolic distances between x and $g(x)$ for $x \in \mathbb{S}_+$. (This equals the length of the closed geodesic represented by g.)

We remark that the positivity of the diffused Margulis-invariants is equivalent to the proper discontinuity of the action of Γ as it is the main result of [28].

3.2. **Neutralized sections.** We note that Γ acts on $\mathbb{US}_+$ as a deck transformation group $\mathbb{U}\Sigma_+$. (We denote $\mathbb{US}_+$ with $\widehat{\mathbb{U}\Sigma_+}$ sometimes and let $\Pi_{\mathbb{U}}$ denote the covering map.) They in [27] following the idea of David Fried constructed a flat affine bundle $\mathbf{E}$ over the unit tangent bundle $\mathbb{U}\Sigma_+$ of Σ_+ by forming $\mathsf{E}^{2,1} \times \mathbb{US}_+$ and taking the quotient by the diagonal action given by the inclusion map

$$h : \Gamma \rightarrow \mathsf{Isom}^+(\mathsf{E}^{2,1}) \subset \mathsf{Aut}(\mathbf{S}^3)$$

where $\gamma(x, v) = (h(\gamma)(x), \gamma(v))$ for a deck transformation γ of the cover $\mathbb{US}_+$ of $\mathbb{U}\Sigma_+$. The cover of $\mathbf{E}$ is denoted by $\hat{\mathbf{E}}$ and is identical with $\mathsf{E}^{2,1} \times \mathbb{US}_+$.

We denote by

$$\pi_{\mathsf{E}^{2,1}} : \hat{\mathbf{E}} = \mathsf{E}^{2,1} \times \mathbb{US}_+ \to \mathsf{E}^{2,1}$$

the projection.

We can form a flat vector bundle $\mathbf{V}$ over $\mathbb{U}\Sigma_+$ by forming $\mathsf{V}^{2,1} \times \mathbb{US}_+$ and taking quotient by the diagonal action given by $\mathcal{L} \circ h$.

A *neutral section* of $\mathbf{V}$ is a $\mathsf{SO}(2,1)$-invariant section which is parallel along geodesic flow of $\mathbb{U}\Sigma_+$. A neutral section $\boldsymbol{\nu} : \mathbb{U}\Sigma_+ \to \mathbf{V}$ arises from a graph of the $\mathsf{SO}(2,1)$-invariant map

$$\tilde{\boldsymbol{\nu}} : \mathbb{US}_+ \to \mathsf{V}^{2,1}$$

with the image in the space of unit space-like vectors in $\mathsf{V}^{2,1}$: $\tilde{\boldsymbol{\nu}}$ is defined by sending a unit vector u in $\mathbb{US}_+$ to the normalization of the Lorentzian cross-product $\rho(\mathsf{u}) \otimes \alpha(\mathsf{u})$ of the null vectors $\rho(\mathsf{u})$ and $\alpha(\mathsf{u})$ respectively representing the the starting point and the ending point in $\mathrm{bd}\mathbb{S}_+$ of the hyperbolic geodesic tangent to u in $\mathbb{S}_+$. (See Section 4.2 of [28] for details.)

Lemma 9. *Suppose that* u *is a unit tangent vector to a geodesic where a deck transformation* γ *acts on along the orientation given by* u*. Then*

- $\rho(\mathsf{u})$ *and* $\alpha(\mathsf{u})$ *respectively correspond in* $\mathbb{S}_+$ *a repelling fixed point and an attracting fixed point on* $\mathrm{bd}\mathbb{S}_+$*,*
- $\tilde{\boldsymbol{\nu}}(\mathsf{u})$ *is an eigenvector of* $\mathcal{L}(\gamma)$ *and corresponds to the saddle-type fixed point of* γ *in* $\mathbb{S}_0$*.*
- γ *acts on a unique space-like line in* $\mathsf{E}^{2,1}$ *in the direction of* $\tilde{\boldsymbol{\nu}}(\mathsf{u})$*, actually by the vector* $\mu(\gamma)\tilde{\boldsymbol{\nu}}(\mathsf{u})$*.*

Proof. Since the geodesic is invariant, the endpoints are attracting or repelling fixed points. The Lorentzian cross-product of the attracting eigenvector and the repelling eigenvector is clearly another eigenvector. The final item is from an interpretation of Margulis invariants. □

Let $\mathbb{U}_{\mathrm{rec}}\Sigma_+$ denote the set of unit tangent vectors of weakly recurrent geodesics on Σ_+. We recall

Lemma 10 ([28])**.** *Let* Σ_+ *be as above. Then*

- $\mathbb{U}_{\mathrm{rec}}\Sigma_+ \subset \mathbb{U}\Sigma_+$ *is a connected compact geodesic flow invariant set and is a subset of the compact set* $\mathbb{U}\Sigma''_+$*.*
- *The inverse image* $\mathbb{U}_{\mathrm{rec}}\mathbb{S}_+$ *of* $\mathbb{U}_{\mathrm{rec}}\Sigma_+$ *in* $\mathbb{U}_{\mathrm{rec}}\mathbb{S}_+$ *is precisely the set of geodesics with both endpoints in* Λ*.*

Proof. The first item is in Lemma 1.2 in [28] and the second item is in the proof of the same lemma. The connectedness is also proved in Lemma 1.3 in [28]. □ □

Let ∇ be the flat connection on $\mathbf{E}$ as a bundle over $\mathbb{US}_+$ with fiber isomorphic to $\mathsf{E}^{2,1}$ induced from the product structure of $\tilde{\mathbf{E}} = \mathsf{E}^{2,1} \times \mathbb{US}_+$. (See Sections 3.2 and 3.3 of [28].)

Using the Anosov geodesic flow of $\mathbf{E}$ in Lemma 8.4 of [28], Goldman, Labourie, and Margulis show that there exists a *neutralized section*, i.e., a section $\mathcal{N} : \mathbb{U}_{\mathrm{rec}}\Sigma_+ \to \mathbf{E}$ satisfying

$$\nabla_X \mathcal{N} = f\boldsymbol{\nu} \tag{6}$$

where X is the vector field of geodesic flow on $\mathbb{U}_{\mathrm{rec}}\Sigma_+$ and f is a real valued function defined on $\mathbb{U}_{\mathrm{rec}}\Sigma_+$. Lemma 3 of [27] shows that we can choose $\mathcal{N}$ so that $f > 0$ by changing $\mathcal{N}$ to $\mathcal{N} + g\boldsymbol{\nu}$ for a suitable real valued $g : \mathbb{U}_{\mathrm{rec}}\Sigma_+ \to \mathbb{R}$.

3.3. **Lifting the neutralized section to the coverings.** Let $\mathbb{U}_{\mathrm{rec}}\mathbb{S}_+$ denote the inverse image of $\mathbb{U}_{\mathrm{rec}}\Sigma_+$ in $\mathbb{U}\mathbb{S}_+$. We take a compact fundamental domain F in $\mathbb{U}_{\mathrm{rec}}\mathbb{S}_+$ of Γ, and then $\mathcal{N}$ lifts on F to a map $\tilde{\mathcal{N}}|F$ to $\tilde{\mathbf{E}}$ so that for each side pairing $g_S \in \Gamma$ of F for a side S of F, we have

$$\begin{aligned} \tilde{\mathcal{N}} \circ g_S(\vec{u}) &= g_S \circ \tilde{\mathcal{N}}(\vec{u}) \\ &= (h(g_S)(\pi_{\mathsf{E}^{2,1}} \circ \tilde{\mathcal{N}}(\vec{u})), g_S(\vec{u})) \text{ for every } \vec{u} \in F. \end{aligned} \tag{7}$$

since $\mathcal{N}$ is obtained from an arbitrary lift satisfying this equation and deforming it (see the proof of Lemma 8.3 of [28]). Thus, we find the section $\tilde{\mathcal{N}} : \mathbb{U}_{\mathrm{rec}}\mathbb{S}_+ \to \tilde{\mathbf{E}}$ lifting $\mathcal{N}$ satisfying

$$\tilde{\mathcal{N}} \circ \gamma = \gamma \circ \tilde{\mathcal{N}} \tag{8}$$

for each deck transformation γ of $\mathbb{U}\mathbb{S}_+ \to \mathbb{U}\Sigma_+$.

Let u' be a unit tangent vector to a closed geodesic in Σ_+. The geodesic flow line α on $\mathbb{U}_{\mathrm{rec}}\Sigma_+$ tangent to u' will return to u' after a period. Suppose that a deck transformation γ in $\mathbb{U}_{\mathrm{rec}}\mathbb{S}_+$ corresponds to a closed geodesic in Σ_+, and acts on a geodesic l_γ in $\mathbb{S}_+$ containing a unit vector u. Let $\overrightarrow{l}_\gamma$ denote the subspace of unit tangent vectors to l_γ in the γ-direction. Since $\tilde{\mathcal{N}}$ is a lift of $\mathcal{N}$, we have that $\pi_{\mathsf{E}^{2,1}} \circ \tilde{\mathcal{N}}|\overrightarrow{l}_\gamma$ maps to a line parallel to $\tilde{\boldsymbol{\nu}}(\mathsf{u})$ up to an action of $\mathcal{L}(\Gamma)$ by equation 6. Since γ acts on the image of this line in $\mathsf{E}^{2,1}$ by equation 8, and $\mathcal{L}(\gamma)$ has $\tilde{\boldsymbol{\nu}}(\mathsf{u})$ as an eigenvector by Lemma 9, we obtain that $\pi_{\mathsf{E}^{2,1}} \circ \tilde{\mathcal{N}}|\overrightarrow{l}_\gamma$ is a line parallel to $\tilde{\boldsymbol{\nu}}(\mathsf{u})$.

Equation 6 also lifts to

$$\nabla_X \tilde{\mathcal{N}} = f\tilde{\boldsymbol{\nu}} \tag{9}$$

where X is the unit-vector field of geodesic flow on $\mathbb{U}\mathbb{S}_+$, ∇ is the flat connection on $\mathsf{E}^{2,1} \times \mathbb{U}\mathbb{S}_+$, and $f : \mathbb{U}_{\mathrm{rec}}\mathbb{S}_+$ is a positive-valued function. Thus, we conclude that the geodesic flow on $\mathbb{U}_{\mathrm{rec}}\mathbb{S}_+$ is conjugated to one on the image subset of $\mathbb{U}\mathsf{E}^{2,1}$ by a map associated with $\pi_{\mathsf{E}^{2,1}} \circ \tilde{\mathcal{N}}$. (We note here that the flow speed changes by the diffused Margulis invariants.)

3.4. **A bounded set weakly recurrent geodesics of Σ_+ goes to a bounded set of ones of $\mathsf{E}^{2,1}/\Gamma$.** Theorem 1 of [27] shows that the image of a geodesic γ of $\mathbb{S}_+$ mapping to a weakly recurrent geodesic in Σ_+ under $\pi_{\mathsf{E}^{2,1}} \circ \tilde{\mathcal{N}}$ is a space-like geodesic in $\mathsf{E}^{2,1}$ mapping to a weakly recurrent geodesic in $\mathsf{E}^{2,1}/\Gamma$.

Let $\mathbb{U}\mathsf{E}^{2,1}$ denote the space of unit space-like vectors at each point of $\mathsf{E}^{2,1}$ and $\mathbb{U}(\mathsf{E}^{2,1}/\Gamma)$ the space of unit space-like vectors at each point of $\mathsf{E}^{2,1}/\Gamma$. Denote by $\mathbb{U}_{\mathrm{rec}}\mathsf{E}^{2,1}$ in $\mathbb{U}\mathsf{E}^{2,1}$ the inverse image of the subset of $\mathbb{U}_{\mathrm{rec}}(\mathsf{E}^{2,1}/\Gamma)$ composed of unit tangent space-like vectors tangent to weakly recurrent space-like geodesics.

We give the topology of the space $\mathcal{G}_{\mathrm{rec}}\mathbb{S}_+$ of oriented geodesics in $\mathbb{S}_+$ mapping to weakly recurrent oriented geodesics in Σ_+ by the quotient topology from the unit tangent bundle $\mathbb{U}_{\mathrm{rec}}\mathbb{S}_+$. Similarly, we give the topology of the space $\mathcal{G}_{\mathrm{rec}}\mathsf{E}^{2,1}$ of space-like oriented geodesics in $\mathsf{E}^{2,1}$ mapping to weakly recurrent geodesics in $\mathsf{E}^{2,1}/\Gamma$ by the quotient topology from the space $\mathbb{U}_{\mathrm{rec}}\mathsf{E}^{2,1}$.

A *bounded subset* of $\mathcal{G}_{\mathrm{rec}}\mathbb{S}_+$ is a set of geodesics passing a bounded set in $\mathbb{U}\mathbb{S}_+$, and a *bounded subset* of $\mathcal{G}_{\mathrm{rec}}\mathsf{E}^{2,1}$ is a set of space-like geodesics passing a bounded set in $\mathbb{U}\mathsf{E}^{2,1}$ under the Euclidean metric d_E.

Proposition 11. *The map $\tilde{\mathscr{N}}$ induces a continuous function $\mathscr{N} : \mathcal{G}_{rec}\mathbb{S}_+ \to \mathcal{G}_{rec}\mathsf{E}^{2,1}$ where*

- *if the oriented geodesic l in $\mathbb{S}_+$ is g-invariant for $g \in \Gamma$, then g acts on the space-like geodesic L_g the image under $\mathscr{N}$ as a translation.*
- *the bounded set of elements of $\mathcal{G}_{rec}\mathbb{S}_+$ maps to a bounded set in $\mathcal{G}_{rec}\mathsf{E}^{2,1}$.*
- *Finally, the map is surjective.*

Proof. For a flow segment in $\mathbb{U}_{\mathrm{rec}}\mathbb{S}_+$, $\pi_{\mathsf{E}^{2,1}} \circ \tilde{\mathscr{N}}$ sends it to a space-like line segment in $\mathsf{E}^{2,1}$. This induces $\mathscr{N}$. The first item is discussed in Section 3.3.

The next statement follows by the fact that $\mathscr{N}$ is induced by the continuous map $\pi_{\mathsf{E}^{2,1}} \circ \tilde{\mathscr{N}}$ which sends a compact set to a compact set.

Clearly our map is surjective from the set of geodesics in $\mathbb{S}_+$ mapping to closed ones in $\mathbb{S}_+/\Gamma$ to the set of geodesics in $\mathsf{E}^{2,1}$ mapping to closed ones in $\mathsf{E}^{2,1}/\Gamma$ since any a closed geodesic correspond to a unique nontrivial element of Γ in the one-to-one manner. The surjectivity of $\mathscr{N}$ now follows by Theorem 1 of [27] by the density of closed geodesics and the continuity of $\mathscr{N}$. □ □

4. THE PROPER-DISCONTINUITY OF THE Γ-ACTION ON $\mathsf{E}^{2,1} \cup \tilde{\Sigma}$.

Let Γ be an orientation-preserving Lorentzian isometry group without any parabolic elements acting freely and properly discontinuously on $\mathsf{E}^{2,1}$ isomorphic to a free group of finite rank ≥ 2. Recall that $\mathbb{S}_+/\Gamma$ is a complete genus $\tilde{\mathsf{g}}$ hyperbolic surface with b ideal boundary components. As above, we assume that the Margulis invariants of nonidentity elements are all positive.

4.1. The action of quasi-Anosov automorphisms.

A homogeneous coordinate system of $\mathbf{S}^3$ is given by setting a point p of $\mathbf{S}^3$ to have coordinates $[t, x, y, z]$ where (t, x, y, z) divided by its norm represent p as a unit vector in $\mathbf{S}^3$. We identified $\mathsf{E}^{2,1}$ with the open affine space given by $t > 0$. We use the coordinates so that $\mathbb{S}$ corresponds to the hyperplane given by $t = 0$ and the origin O of $\mathsf{E}^{2,1}$ is given by $[1, 0, 0, 0]$. The closure of $\mathsf{E}^{2,1}$ is the hemisphere

$\mathscr{H}$ with $\partial\mathscr{H} = \mathbb{S}$. For this system of coordinates, $(x, y, z) \in \mathsf{E}^{2,1}$ is given homogeneous coordinates $[1, x, y, z]$.

A projective automorphism g that is of form

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \lambda & 0 & 0 \\ k & 0 & 1 & 0 \\ 0 & 0 & 0 & \frac{1}{\lambda} \end{bmatrix} \lambda > 1, k \neq 0 \tag{10}$$

under a homogeneous coordinate system of $\mathbf{S}^3$ is said to be a *quasi-Anosov projective automorphism*, generalizing the terminology "Anosov transformation". To explain the terminology, g acts on an open hemisphere to be identified with $\mathsf{E}^{2,1}$ and has an invariant line given by $x = 0, z = 0$ where it acts as a translation and the invariant plane S given by $x = 0$ and one U given by $z = 0$ containing it. On S, g contracts vectors along a direction by a constant $1/\lambda$ and g expands the vectors along a direction by a constant λ on U assuming $\lambda > 1$. The plane S given by $x = 0$ is said to be the *stable plane*, and the plane U given by $z = 0$ is said to be the *unstable plane*.

Obviously a Lorentzian isometry with a nontrivial translational part is quasi-Anosov.

Let η_+ be the segment with endpoints $[0, 1, 0, 0]$, $[0, -1, 0, 0]$ passing $[0, 0, 1, 0]$ in the middle. Let η_- denote the segment with endpoints $[0, 0, 0, 1]$ and $[0, 0, 0, -1]$ passing $[0, 0, -1, 0]$. These have $\mathbf{d}_{\mathbf{S}^3}$-lengths $= \pi$.

Lemma 12 (Central). *Let $g_{\lambda,k}$ denote the automorphism on $\mathbf{S}^3$ defined by the above matrix in the equation 10 for a homogeneous coordinate system with coordinate functions t, x, y, z in the given order and let $\mathbf{S}^2_0$ denote the subspace given by $x = 0$. We assume that $k \geq 0, \lambda > 0$. Then as $\lambda, k \to +\infty$ where $k/\lambda \to 0$, we obtain*

- *$g_{\lambda,k}|\mathbf{S}^3 - \mathbf{S}^2_0$ converges in the compact open topology to a rational map Π_0 given by sending $[t, x, y, z]$ to $[0, \pm 1, 0, 0]$ where the sign depends on the sign of x/t if $t \neq 0$ and the sign of x if $t = 0$.*
- *On $g_{\lambda,k}|(\mathbf{S}^2_0 \cap \mathscr{H}) - \eta_-$ converges in the compact open topology to a rational map Π_1 given by sending $[t, 0, y, z]$ to $[0, 0, 1, 0]$.*
- *For a properly convex compact set K in $\mathscr{H} - \eta_-$, the geometric limit of a subsequence of $\{g_{\lambda,k}(K)\}$ as $\lambda, k \to \infty$, is either a point $[0, 1, 0, 0]$ or $[0, -1, 0, 0]$ or the segment η_+.*

Proof. The first two items follow by normalizing the above matrix by diving it by the largest entry.

For any properly convex compact subset K' in $\mathscr{H} - \mathbf{S}^2_0$, $\{g_{\lambda,k}(K')\}$ converges to one of $\{[0, 1, 0, 0]\}$ or $\{[0, -1, 0, 0]\}$ as $\lambda, k \to \infty$ while $k/\lambda \to 0$ by the first item.

So, we now suppose that $K' \cap \mathbf{S}^2_0 \neq \emptyset$. For a properly convex compact subset K' in $\mathscr{H} - \eta_-$, let K'_1 be the nonempty properly convex set $K' \cap \mathbf{S}^2_0$. We can take an open 2-hemisphere H with boundary $\mathrm{bd}H \subset \mathbb{S}$ so that H is the topological boundary in $\mathscr{H}$ of a convex open domain $H' \subset \mathscr{H}$



FIGURE 5. The action of a Lorentzian isometry $\hat{g}$ on the hemisphere $\mathscr{H}$ where the boundary sphere $\mathbb{S}$ is the unit sphere with center $(0,0,0)$ here. We are using different homogeneous coordinates. The arc on $\mathbb{S}$ given by $y=0$ is the invariant geodesic in $\mathbb{S}_+$ and with end points the fixed points of $\hat{g}$. The arc given by $x=0$ and $z=0$ is a line where $\hat{g}$ acts as a translation in the positive y-axis direction for $\hat{g}\neq \mathrm{I}$. The plane $z=0$ is where $\hat{g}$ acts as an expansion-translation and the plane $x=0$ is where $\hat{g}$ acts as a contraction-translation. The semicircle defined by $y\geq 0$ and $z=0$ is η^+, "the attracting arc". The semicircle defined by $x=0$ and $y\leq 0$ is η^-, "the repelling arc".

disjoint from η_- and containing K' where we can freely and slightly enlarge subspaces and take joins. (For example, we can choose H to be a plane in $\mathsf{E}^{2,1}$ separating K' and η_- containing $[0,0,0,\pm 1]$ in the boundary.) Then the following statements hold:

- there exist a properly convex compact domain D_1 in a component of $H'-\mathbf{S}^2_0$ with $x>0$ and another one D_2 in a component of $H'-\mathbf{S}^2_0$ with $x<0$ so that K' is in the join J of D_1 and D_2
- For $K_1 := J\cap \mathbf{S}^2_0 \supset K'_1$, the join J_1 of K_1 and D_1 and the join J_2 of K_1 and D_2 satisfy $J = J_1 \cup J_2$.

For example, we may obtain D_1 and D_2 by removing from K' an ϵ-$\mathbf{d}_{\mathbf{S}^3}$-neighborhood of $\mathbf{S}^2_0$ for small $\epsilon > 0$ and making them slightly larger so that the join of D_1 and D_2 contains K'.

By the second item of the lemma that we are proving currently, we obtain that $\{g_{\lambda,k}(K_1)\}$ converges to $[0,1,0,0]$ as $\lambda, k \to \infty$. Since $\{g_{\lambda,k}(D_1)\} \to \{[0,1,0,0]\}$ and $\{g_{\lambda,k}(D_2)\} \to \{[0,-1,0,0]\}$ as $\lambda, k \to \infty$ by above, we see that

- $\{g_{\lambda,k}(J_1)\}$ geometrically converge to the unique convex segment s_1 with endpoints $[0,1,0,0]$ and $[0,0,1,0]$ in $\mathbb{S}$ and
- $\{g_{\lambda,k}(J_2)\}$ geometrically converge to the unique convex segment s_2 with endpoints $[0,-1,0,0]$ and $[0,0,1,0]$ in $\mathbb{S}$ by Lemma 4 respectively.

Hence, we conclude $\{g_{\lambda,k}(J)\} \to \eta_+$ as $\lambda, k \to \infty$.

For each $\epsilon > 0$, we have that

$$g_{\lambda,k}(K') \subset N_\epsilon(\eta_+) = N_\epsilon(s_1 \cup s_2)$$

whenever λ and k are sufficiently large, and there are points arbitrarily close to both points $[0,\pm1,0,0]$. As K' is convex and connected, the geometric limit of $\{g_{\lambda,k}(K')\}$ exists as $\lambda, k \to \infty$ and equals η_+. (See Figure 5 also.)

□ □

We remark that our proof of Proposition 13 is a generalization of the above proof.

4.2. **The proper discontinuity of the action of Γ.** Recall from Theorem 8 that we can construct

$$\tilde{\Sigma} := \mathbb{S}_+ \cup \coprod_{i \in \mathcal{J}} \mathcal{R}_i \cup \mathbb{S}_- \subset \mathbb{S}$$

that is a Γ-invariant open domain with an infinite fundamental group and covers a closed real projective surface Σ, a union of Σ'_+ and Σ'_- and annuli A_i, $i = 1, \dots, \mathsf{b}$.

Proposition 13. *Let Γ be a discrete group of orientation-preserving Lorentzian isometries acting freely and properly discontinuously on $\mathsf{E}^{2,1}$ isomorphic to a free group of finite rank ≥ 2 with $\tilde{\Sigma}$ as determined above. Then Γ acts freely and properly discontinuously on $\mathsf{E}^{2,1} \cup \tilde{\Sigma}$ as a group of projective automorphisms of $\mathbf{S}^3$.*

Proof. Since Γ acts freely and properly discontinuously on $\mathbb{R}^3 = \mathsf{E}^{2,1}$ and so on the constructed $\tilde{\Sigma}$ separately, Γ acts freely on the union.

As we said before, we assume without loss of generality that Margulis invariants of nonidentity elements are all positive. (If the signs are all negative, then actually, we need to choose $\tilde{\Sigma}_-$ instead of $\tilde{\Sigma}$.)

Suppose that there exists a sequence $\{g_i\}$ of elements of Γ and a compact subset K of $\mathsf{E}^{2,1} \cup \tilde{\Sigma}$ so that

$$g_i(K) \cap K \neq \emptyset \text{ for all } i. \tag{11}$$

We will show that actually the sequence $\{g_i\}_{i\in\mathbb{Z}^+}$ has to be a finite sequence for the set of positive integers $\mathbb{Z}^+$.

We prove by contradiction: Suppose that $\{g_i\}$ is an infinite sequence and $g_i \neq g_j$ if $i \neq j$. The equation 11 holds universally in this section even if we take any subsequences.

Recall that the Fuchsian Γ-action on the boundary $\mathrm{bd}\mathbb{S}_+$ of the standard disk $\mathbb{S}_+$ in $\mathbb{S}$ forms a discrete convergence group: For every sequence g_j in Γ, there is a subsequence g_{j_k} and two (not necessarily distinct) points a, b in the circle $\mathrm{bd}\mathbb{S}_+$ such that

- the sequences $g_{j_k}(x) \to a$ locally uniformly in $\mathrm{bd}\mathbb{S}_+ - \{b\}$, i.e., uniformly in any compact subset of $\mathrm{bd}\mathbb{S}_+ - \{b\}$ and
- $g_{j_k}^{-1}(y) \to b$ locally uniformly on $\mathrm{bd}\mathbb{S}_+ - \{a\}$ respectively as $k \to \infty$.

(See [1] for details.) We remark that

$$a, b \in \Lambda. \tag{12}$$

(I) *First, we will consider the case when* $a \neq b$:

(I)(i) *We collect some boundedness properties of* [28]: Now letting $\{g_i\}$ denote the subsequence, we note that each g_i has an attracting fixed point a_i and a repelling fixed point r_i in $\mathrm{bd}\mathbb{S}_+$. By our conditions, sequences $\{a_i\} \to a$ and $\{r_i\} \to b$.

A Lorentzian isometry element g_i acts as a translation on a unique space-like line L_{g_i} in the direction ν_i of eigenvalue 1. Let α_i and ρ_i denote the null vectors in the directions of a_i and r_i respectively. We compute

$$\nu_i := \frac{\rho_i \otimes \alpha_i}{|||\rho_i \otimes \alpha_i|||}$$

i.e., the Lorentzian cross-product of ρ_i and α_i representing r_i and a_i respectively divided by the absolute value of its norm. (We consider r_i as the start and a_i as the ending point and ν_i as the neutral vector.) Since we have $\{a_i\} \to a$, we obtain that the sequence $\overline{a_i[\nu_i]a_{i,-}} = \mathrm{Cl}(\varepsilon(a_i))$ converges to a segment $\overline{a[\nu]a_-} = \mathrm{Cl}(\varepsilon(a))$ where $[\nu]$ is the direction of

$$\nu := \frac{\beta \otimes \alpha}{|||\beta \otimes \alpha|||}$$

the Lorentzian cross-product of nonzero vectors α and β corresponding to a and b respectively.

Since the geodesics with end points a_i, r_i pass the bounded part of the unit tangent bundle of $\mathbb{S}_+$, it follows that L_{g_i} are bounded as well by Proposition 11. Each L_{g_i} pass a point p_i, and $\{p_i\}$ forms a bounded sequence in $\mathsf{E}^{2,1}$. By choosing a subsequence, we assume without loss of generality that $p_i \to p_\infty$ for $p_\infty \in \mathsf{E}^{2,1}$

(I)(ii) *The next important step of the proof is that the coordinate changes required for* g_i *so that it becomes of form in equation* 10 *are uniformly bounded*:

We can conjugate by a sequence of h_i of orientation-preserving quasi-isometries considered as an element of $\mathsf{Aut}(\mathbf{S}^3)$ so that $h_i g_i h_i^{-1}$ is of form

$$\hat{g}_i := \left[\begin{array}{cccc} 1 & 0 & 0 & 0 \\ 0 & \lambda(g_i) & 0 & 0 \\ k(g_i) & 0 & 1 & 0 \\ 0 & 0 & 0 & \frac{1}{\lambda(g_i)} \end{array}\right]. \tag{13}$$

where $\lambda(g_i) > 1, k(g_i) > 0$ and the orientation preserving h_i is given by sending $[0,1,0,0]$ to a_i and $[0,0,0,1]$ to r_i and $[1,0,0,0]$ to p_i and $[0,0,1,0]$ to $[\nu_i]$. (We choose h_i to acts on $\mathscr{H}$ and hence an affine transformation.) Here the positivity of the Margulis invariant is equivalent to $k(g_i) > 0$ as the coordinate change is orientation-preserving. (Note here that $k(g_i)$ depends on the coordinate system.)

We note that

$$g_i = h_i^{-1} \circ \hat{g}_i \circ h_i; \tag{14}$$

hence, g_i acts like the standard form matrix in equation 13 up to coordinatization by h_i.

Since

- the coordinate axes are associated with three eigenvectors α_i, ν_i, ρ_i where ν_i is of Lorentzian unit-norm and is the Lorentzian cross-product of ρ_i, α_i where $a_i = [\alpha_i], r_i = [\rho_i]$.
- $a_i \to a$ and $r_i \to b$, $\rho_i \otimes \alpha_i = \nu_i \to \nu$, and
- $p_i \to p$,

we assume without loss of generality by choosing a subsequence that h_i converges to a quasi-isometry h uniformly in C^∞-sense. Hence the sequence $\{h_i\}$ is *uniformly quasi-isometric* in $\mathbf{d}_{\mathbf{S}^3}$; that is,

$$C^{-1}\mathbf{d}_{\mathbf{S}^3}(x,y) \leq \mathbf{d}_{\mathbf{S}^3}(h_i(x), h_i(y)) \leq C\mathbf{d}_{\mathbf{S}^3}(x,y) \text{ for } x, y \in \mathbf{S}^3, i \in \mathbb{Z}_+ \tag{15}$$

for a fixed $C > 0$ independent of i by Proposition 11.

The ith-coordinate system is a coordinate system where we have

$$\begin{aligned} h_i(p_i) &= [1,0,0,0], h_i(a_i) = [0,1,0,0], \\ h_i(b_i) &= [0,0,0,1], \text{ and } h_i([\nu_i]) = [0,0,1,0]. \end{aligned} \tag{16}$$

Moreover, L_{g_i} is sent to

$$\overline{[0,0,1,0][1,0,0,0],[0,0,-1,0]}^o.$$

The sequence of each kind of ith-coordinate functions converges to a coordinate function of a new adopted coordinate system h where we have

$$\begin{aligned} h(p) &= [1,0,0,0], h(a) = [0,1,0,0], \\ h(b) &= [0,0,0,1], \text{ and } h([\nu]) = [0,0,1,0]. \end{aligned} \tag{17}$$

Moreover, at the same time, we can assume that the linearization of the affine maps h_i in $\mathsf{E}^{2,1}$ sends the vector ν_i to $(0,1,0)$ in $\mathsf{V}^{2,1}$ and, similarly, h sends the vector ν to $(0,1,0)$.

We recall from Lemma 12:

- the subspace $\mathbf{S}^2_0$ given by $x = 0$,
- the segment η_+ with endpoints $[0, 1, 0, 0]$, $[0, -1, 0, 0]$ passing $[0, 0, 1, 0]$ in the middle, and
- the segment η_- with endpoints $[0, 0, 0, 1]$ and $[0, 0, 0, -1]$ and passing $[0, 0, -1, 0]$.

Then

- $h_i^{-1}(\eta_-) = \mathrm{Cl}(\varepsilon(r_i))$ is a segment of $\mathbf{d}_{\mathbf{S}^3}$-length π with endpoints $r_i, -r_i$ and passing $[-\nu_i]$ and
- $h_i^{-1}(\eta_+) = \mathrm{Cl}(\varepsilon(a_i))$ is a segment of $\mathbf{d}_{\mathbf{S}^3}$-length π with endpoints $a_i, -a_i$ and passing $[\nu_i]$.

Finally for this step (I)(ii), *we estimate the matrix entries*: Using the hyperbolic metric on $\mathbb{S}_+$, we see that the closed geodesic c_i in Σ_+ represented by g_i has the hyperbolic length

$$l_{\mathbb{S}_+}(g_i) = \left|\log \frac{\lambda(g_i)}{\frac{1}{\lambda(g_i)}}\right| = 2|\log \lambda(g_i)|,$$

where we used cross ratios.

Since g_i is divergent as it acts on $\mathbb{S}_+$, we have that

Lemma 14 (Choi, Goldman)**.** *By conjugating g_i by h_i as defined above to the form of equation 13, we have*

$$\lambda(g_i) \to +\infty, \text{ and } k(g_i) \to +\infty, k(g_i)/\lambda(g_i) \to 0. \tag{18}$$

Proof. Since $\{g_i\}$ restricts to an unbounded sequence hyperbolic isometries in $\mathbb{S}_+$, it follows that $\{\lambda(g_i)\} \to \infty$.

We claim that

$$\mu(g_i) = k(g_i): \tag{19}$$

we chose x in $\mathbf{B}(g_i(x) - x, v_0(g_i))$ to be on L_{g_i} and the linearization $\mathcal{L}(h_i)$ of h_i restricted to $\mathsf{E}^{2,1}$ sends $g_i(x) - x$ and $\mathsf{v}_0(g_i)$ respectively to a vector parallel to $(0, 1, 0)$ and the vector $(0, 1, 0)$. Since $||\mathsf{v}_0(g_i)|| = 1$, we obtain $g_i(x) - x = \mu(g_i)\mathsf{v}_0(g_i)$. Hence, $\mathcal{L}(h_i)(g_i(x) - x) = \mu(g_i)(0, 1, 0)$. Since $k(g_i) = \mathcal{L}(h_i)(g_i(x) - x) \cdot (0, 1, 0)$, we have the equality 19. By equation 5, we have $\mu(g_i) \to \infty$. Hence, the second limit follows.

By the main corollary of [28], we have $\mu(g_i) \leq C'l(g_i)$ and hence $\mu(g_i) \leq 2C'|\log \lambda(g_i)|$; we have the final limit. $\square$ $\square$

(I)(iii) *We go back to our original compact set K from the beginning of the proof and cover it by compact convex sets that we can understand under the action of g_i*:

Recall from (I)(ii) that we may assume by choosing subsequences of g_i that

- $\{h_i^{-1}(\mathbf{S}^2_0)\} \to \mathbf{S}^2_\infty$,
- $\{h_i^{-1}(\eta_-)\} \to \eta_-^\infty = \mathrm{Cl}(\varepsilon(b))$, and
- $\{h_i^{-1}(\eta_+)\} \to \eta_+^\infty = \mathrm{Cl}(\varepsilon(a))$.

Here

- η_+^∞ is a segment with endpoints a and a_- passing $[\nu]$ of $\mathbf{d}_{\mathbf{S}^3}$-length π and
- η_-^∞ is a segment with endpoints b and $-b$ passing $[-\nu]$ of $\mathbf{d}_{\mathbf{S}^3}$-length π.

They are all tangent to $\mathrm{bd}\mathbb{S}_+$.

Note that K is a subset of $\mathscr{H} - \bigcup_{i\in\mathbb{Z}_+} h_i^{-1}(\eta_-)$ since for each i, $h_i^{-1}(\eta_-) = \mathrm{Cl}(\varepsilon(r_i))$ is in the complement of $\tilde{\Sigma}$ by definition. We define the compact set $K_i := K \cap h_i^{-1}(\mathbf{S}_0^2)$. Now fix a sufficiently small ϵ, δ so that $0 < \delta < \epsilon$. We cover K

- by compact properly convex balls B_j, $j \in I_1$ for a finite index set I_1, in $\mathsf{E}^{2,1} \cup \tilde{\Sigma}$ that are at least ϵ away from $\mathbf{S}_\infty^2$ by $\mathbf{d}_{\mathbf{S}^3}$-distances and
- by small compact properly convex balls C_j, $j \in I_2$ for another finite index set I_2, which for each j is the join of two properly convex balls C_j^+ and C_j^- in $\mathsf{E}^{2,1} \cup \tilde{\Sigma}$ at least δ away from $\mathbf{S}_\infty^2$ in $\mathbf{d}_{\mathbf{S}^3}$-distances.

This can be achieved by first covering by balls at least ϵ-away from $\mathbf{S}_\infty^2 \cap K$ and balls with centers on $\mathbf{S}_\infty^2 \cap K$ and dividing the slightly larger balls containing the second class of balls into two parts and pushing these away from $\mathbf{S}_\infty^2$.

There are two components H_a and H_{a_-} of $\mathscr{H} - \mathbf{S}_\infty^2$ containing a or a_- respectively. We divide I_1 into I_1^+ and I_1^- so that $j \in I_1^+$ if and only if $B_j \subset H_a$. Also, we assume without loss of generality that $C_j^+ \subset H_a$ and $C_j^- \subset H_{a_-}$ for $j \in I_2$. We denote by $C_{j,i}^0$ for $j \in I_2$, the set $h_i(\mathbf{S}_0^2) \cap J(C_j^+, C_j^-)$.

Since $\{h_i^{-1}(\mathbf{S}_0^2)\}$ converges to $\mathbf{S}_\infty^2$, we have the following:

- There exists I_0 such that for $i > I_0$, $h_i^{-1}(\mathbf{S}_0^2)$ is at least $\delta/8$ away from $\bigcup_{j\in I_1^+} B_j$ and $\bigcup_{j\in I_1^-} B_j$ and from $\bigcup_{j\in I_2} C_j^+$ and $\bigcup_{j\in I_2} C_j^-$.
- For $i > I_0$ and a positive constant C, $\mathbf{S}_0^2$ is at least $C^{-1}\delta/8$ away from $h_i(\bigcup_{j\in I_1^+} B_j)$ and $h_i(\bigcup_{j\in I_1^-} B_j)$ and from $h_i(\bigcup_{j\in I_2} C_j^+)$ and $h_i(\bigcup_{j\in I_2} C_j^-)$ since h_i, $i \in \mathbb{Z}_+$, are uniformly quasi-isometric.
- Now we act by $\hat{g}_i$ and then by h_i^{-1}.
- Hence, for every $\epsilon'' > 0$, we have an integer $I_0' > I_0$ so that for $i > I_0'$, the images of these sets under g_i go into the ϵ''-$\mathbf{d}_{\mathbf{S}^3}$-neighborhoods of a or a_- by Lemma 12 since $\{h_i^{-1}\}$ is uniformly quasi-isometric in $\mathbf{d}_{\mathbf{S}^3}$. (See equations 14 and 15.)

Let us fix $j \in I_2$. We first recall that the sets C_j^0 are all uniformly bounded away from $h_i^{-1}(\eta_-) = \mathrm{Cl}(\varepsilon(r_i))$ for $i > I_0$ for sufficiently large I_0. Hence $h_i(C_j^0)$ are all uniformly bounded away from η_- for $i > I_0$ for sufficiently large I_0 as $\{h_i\}$ is uniformly quasi-isometric. Furthermore, the sequence of images under $\{\hat{g}_i\}$ of C_j^0 for each j then converges to $[0, 0, 1, 0]$ by Lemma 12. Therefore, the sequence $\{g_i(C_j^0)\}_{i\in\mathbb{Z}_+}$ converges to $\{[\nu]\}$ by equation 14 since the collection of automorphisms h_i^{-1} are uniformly quasi-isometric in $\mathbf{d}_{\mathbf{S}^3}$.

Now, the sequence

- $\{g_i(J(C^j_+, C^j_0))\}_{i\in\mathbb{Z}_+} \to \overline{a[\nu]}$ by Lemma 4.
- Similarly, $\{g_i(J(C^j_-, C^j_0))\}_{i\in\mathbb{Z}_+} \to \overline{[\nu]a_-}$.

Since $J(C^j_+, C^j_-) = J(C^j_+, C^j_0) \cup J(C^j_-, C^j_0)$, we obtain that

$$\{g_i(J(C^j_+, C^j_-))\}_{i\in\mathbb{Z}_+} \to \eta^\infty_+ = \overline{a[\nu]} \cup \overline{[\nu]a_-} \text{ for each } j \in I_2.$$

Thus, we see that if $g_i(K)\cap K \neq \emptyset$ for all i, then there has to be some fixed pair of balls B and B' in the above finite collection of balls $I_1 \cup I_2$ so that $g_i(B)\cap B' \neq \emptyset$ for infinitely many i. However, $\{g_i(B)\}$ converges to a subset of $\eta^\infty_+ \subset \mathbb{S}$ with $\eta^\infty_+ \cap \tilde{\Sigma} = \emptyset$. Thus, for each $\epsilon > 0$, there exists an integer I_0 such that $N_\epsilon(\eta^\infty)$ contains $g_i(B)$ if $i > I_0$. However, $N_\epsilon(\eta^\infty_+)\cap B' = \emptyset$ for any ball in $I_1 \cup I_2$ for sufficiently small ϵ, and we obtain a contradiction.

(II) Now we consider the $a = r$ case. Choose an element $\gamma_0 \in \Gamma$ so that $\gamma_0(a) \neq a$ as Γ is not an elementary Fuchsian group acting on $\mathbb{S}_+$.

Consider the sequence $\{\gamma_0 g_i\}$. For each $\epsilon > 0$ and sufficiently large i, a precompact neighborhood U in $\mathrm{bd}\mathbb{S}_+ - \{a\}$ maps into the ϵ-$\mathbf{d}_{\mathbf{S}^3}$-neighborhood of $\gamma_0(a)$ by $\gamma_0 g_i$. Now consider the sequence $g_i^{-1}\gamma_0^{-1}$. Let V be a precompact neighborhood in $\mathrm{bd}\mathbb{S}_+ - \{\gamma_0(a)\}$. Then $\gamma_0^{-1}(V)$ is a precompact interval in $\mathrm{bd}\mathbb{S}_+ - \{a\}$ and for arbitrary $\epsilon > 0$, g_i^{-1} sends it to an ϵ-$\mathbf{d}_{\mathbf{S}^3}$-neighborhood of a for i sufficiently large.

We note that $\gamma_0 g_i(K \cup \gamma_0(K)) \cap (K \cup \gamma_0(K)) \neq \emptyset$ for infinitely many i. Thus, we are reduced to the case where $a \neq r$ by replacing g_i with $\gamma_0 g_i$ and K with another compact set $K \cup \gamma_0(K) \subset \mathsf{E}^{2,1} \cup \tilde{\Sigma}$. Again, we obtain contradiction.

□

Remark 15. We can replace the condition of the proper-discontinuity of the action of Γ in $\mathsf{E}^{2,1}$ by the condition given by equation 5. Then we can show that this condition implies that Γ acts properly discontinuously on $\mathsf{E}^{2,1}$ giving us an alternative proof of a part of the results of [28] that the positiveness of diffused Margulis invariants implies the proper-discontinuity of the action of Γ on $\mathsf{E}^{2,1}$ since we are only using the property given by the equation 5 and not the proper-discontinuity of the action of Γ on $\mathsf{E}^{2,1}$ itself.

5. The proof of tameness.

Using the above notations, we note that $\tilde{\Sigma}/\Gamma$ is a closed surface of genus g and forms the boundary of the 3-manifold $M := (\mathsf{E}^{2,1} \cup \tilde{\Sigma})/\Gamma$ by Proposition 13. We now show that M is compact.

Proposition 16. *Each simple closed curve γ in $\tilde{\Sigma}$ bounds a simple disk in $\mathsf{E}^{2,1} \cup \tilde{\Sigma}$. Let c be a simple closed curve in Σ that is homotopically trivial in M. Then c bounds an imbedded disk in M.*

Proof. This is just Dehn's lemma. □

We can find a collection of disjoint simple curves γ_i, $i \in \mathcal{J}$, on $\tilde{\Sigma}$ for an index set $\mathcal{J}$ so that the following hold:

- $\bigcup_{i\in\mathcal{J}} \gamma_i$ is invariant under Γ.
- $\bigcup_{i\in\mathcal{J}} \gamma_i$ cuts $\tilde{\Sigma}$ into a union of open pair-of-pants P_k, $k \in K$, for an index set K. The closure of each P_k is a closed pair-of-pants.
- $\{P_k\}_{k\in K}$ is a Γ-invariant set.
- Under the covering map $\pi : \tilde{\Sigma} \to \tilde{\Sigma}/\Gamma$, each γ_i for $i \in I$ maps to a simple closed curve in a one-to-one manner and each P_k for $k \in K$ maps to an open pair-of-pants as a homeomorphism.

This is achieved by finding arcs in $\mathbb{S}_+/\Gamma$ cutting it into disks. Recalling that $\mathbb{S}_+/\Gamma$ is an open surface compactified to the compact surface Σ'_+ with boundary, we obtain a system of geodesic segments $\hat{\alpha}_i$, $i \in I$ for a finite set I, Σ'_+ cutting it into a union of disks each of which is bounded by six arcs, alternating triple of which are arcs coming from $\partial\Sigma'_+$ and the other alternating triples are geodesic segments ending $\partial\Sigma'_+$. We can assume that each $\hat{\alpha}_i$ is imbedded in $\mathbb{S}_+/\Gamma$ cutting it into finitely many hexagons so that each ideal boundary component of $\mathbb{S}_+/\Gamma$ meets $\hat{\alpha}_i$ and $\hat{\alpha}_j$ for some pair i, j, $i \neq j$.

This gives a system of geodesic arcs α_j, $j \in J$, in $\mathbb{S}_+$ for some infinite index set J which decomposes $\mathbb{S}_+$ into hexagons and forms a Γ-invariant set. We define $\alpha_{i,-} := -\alpha_i$ in $\mathbb{S}_-$. We connect each endpoint x of α_i with its antipodal endpoint of $\alpha_{i,-}$ by $\varepsilon(x)$. (These form so-called crooked circles.) We do this for each arc and obtain the above system of simple closed curves. By construction, $\{\gamma_i\}_{i\in J}$ maps to a system of disjointly imbedded curves $\hat{\gamma}_1, .., \hat{\gamma}_{3\mathsf{g}-3}$, and each of them has a simple closed lift in $\tilde{\Sigma}$ and hence has a trivial holonomy. See Figure 6. Also, the collection $\hat{\gamma}_1, \dots, \hat{\gamma}_{3\mathsf{g}-3}$ decomposes the closed surface $\tilde{\Sigma}/\Gamma$ of genus g for some $\mathsf{g} \geq 2$ into $2\mathsf{g}-2$ pairs of pants $P'_1, \dots, P'_{2\mathsf{g}-2}$. (Here g depends on the genus $\tilde{\mathsf{g}}$ of $\mathbb{S}_+/\Gamma$ and the number of ideal boundary components.)

By trivial holonomy and Dehn's lemma, each $\hat{\gamma}_i$ bounds a disk D_i in M. By the 3-manifold topology of disk exchanges, we can choose $D_1, \dots, D_{3g-3}$ to be mutually disjoint. (See [29].)

By combinatorics of the situation, the closure of each component of $M - D_1 - \cdots - D_{3g-3}$ is homeomorphic to a compact 3-ball in as it is bounded by a 2-sphere that is a union of a pair-of-pants P'_j and three of $D_1, \dots, D_{3g-3}$: We can lift the sphere into $\mathsf{E}^{2,1} \cup \tilde{\Sigma}$ and being a sphere which can be pushed inside $\mathsf{E}^{2,1}$, it bounds a compact 3-ball in it disjoint from other disks.

The image of the 3-ball is a compact 3-ball in M as well. Since M is a union of the closure of the components that are 3-balls meeting one another in disjoint disks, we conclude that M is a compact 3-manifold. This implies also that M is homeomorphic to a solid handle body of genus g; that is, our manifold $(\mathsf{E}^{2,1} \cup \tilde{\Sigma})/\Gamma$ is a compact 3-manifold and so its interior $\mathsf{E}^{2,1}/\Gamma$ is tame. This completes the proof of Theorem 2. (Also, we can see that the rank of Γ equals g.)



FIGURE 6. The arcs in $\mathbb{S}_+$ and an example of $\hat{\gamma}_i$ in the bold arcs.

A *crooked plane* is a closed disk embedded in $\mathsf{E}^{2,1}$ constructed as follows: Take two null vectors v_1 and v_2 and respective parallel lines L_1 and L_2 meeting at a point x. Then they are on a time-like plane P. Take all time-like lines on P from x and take their union. It is a union of two cones C_1 and C_2 with vertex at x. We obtain two null half-planes $\mathscr{W}(\mathsf{v}_1)$ and $\mathscr{W}(\mathsf{v}_2)$. A crooked plane is the union $C_1 \cup C_2 \cup \mathscr{W}(\mathsf{v}_1) \cup \mathscr{W}(\mathsf{v}_2)$. (See [15] for details.)

A disk in $\mathsf{E}^{2,1}$ is an *almost crooked plane* if it agrees with a crooked plane in the complement of its compact subset. Also, its immersed or imbedded image is said to be *crooked plane* as well.

Proposition 17. *Let* $\tilde{\Sigma}$ *and* Γ *be as above. The Margulis space-time* $\mathsf{E}^{2,1}/\Gamma$ *with a free fundamental group has a system of disks with boundary in* $\mathsf{E}^{2,1} \cup \tilde{\Sigma}/\Gamma$ *so that the closures of components of the complement are compact* 3-*balls and the disks are almost crooked planes.*

Proof. Each of our disks has the boundary of a crooked plane. In the Dehn's lemma, one can arbitrarily assign the tubular neighborhood of the boundary of each disk as long as it is transversal to $\tilde{\Sigma}/\Gamma$. $\square$

6. ASYMPTOTICS OF MARGULIS SPACE-TIMES

Let Γ be as above. We show the existence of two convex hypersurfaces in $\mathsf{E}^{2,1}$ asymptotic to $\mathrm{bd}\mathbb{S}_+$ and $\mathrm{bd}\mathbb{S}_-$ respectively. We will use this to show that a concave boundary submanifold contains all the closed space-like geodesics in $\mathsf{E}^{2,1}/\Gamma$ in the next section.

6.1. **Supporting hyperplanes.** Let $\mathcal{H}$ be the set of pairs (x,h) where x is a point of the limit set Λ in $\mathrm{bd}\mathbb{S}_+$ and a 2-hemisphere h in $\mathscr{H}$ containing x in its boundary ∂h. We give the Hausdorff convergence topology to this space. γ in Γ acts by sending (x,h) to $(\gamma(x),\gamma(h))$.

Proposition 18. *For each point x of the limit set Λ of Γ in $\mathbb{S}$ in* $\mathrm{bd}\mathbb{S}_+$*, there exists a unique 2-dimensional hemisphere h_x depending continuously on x so that h_x contains the fixed line $L_g \subset \mathbb{R}^3$ of an element of $g \in \Gamma$ if x is an attracting fixed point of g. Moreover, there exists a map $\tau : \Lambda \to \mathcal{H}$ sending x to (x,h_x) that is continuous and the map is Γ-equivariant; i.e.,*

$$\tau(\gamma(x)) = (\gamma(x),\gamma(h_x)) \text{ for } \tau(x) = (x,h_x).$$

Proof. If x is an attracting fixed point of $\gamma \in \Gamma$, we let h_x be the hemisphere containing the invariant line $L_\gamma \subset \mathsf{E}^{2,1}$ of γ and x as the stabilizer subgroup of Γ is infinite cyclic and is uniquely determined by x by the elementary Fuchsian group theory. Then h_x is γ-equivariant and ∂h_x contains $x, -x, s, -s$ in $\mathbb{S}$ for the saddle-type fixed points s and $-s$ and the antipode $-x$ of x. Thus, τ is defined for the set of attracting fixed points of Γ.

The inverse image in $\mathbb{U}\mathbb{S}_+$ of $\mathbb{U}_{\mathrm{rec}}(\mathbb{S}_+/\Gamma)$ is precisely the set of geodesics with endpoints in Λ. (See Lemma 10 and the proof of Lemma 1.3 of [28].) Recall that $\Lambda^* = \Lambda \times \Lambda - \Delta$ consists of elements (x,y) where x is the ending point and y is the starting point of a weakly recurrent geodesic l.

Denote by $\overline{xy}^o$ the open line segment connecting two points $x,y \in \mathrm{bd}\mathbb{S}_+$. For convenience, we use the notation from Proposition 11

$$L(x,y) := \mathscr{N}(\overline{xy}^o).$$

Using Proposition 11, we define a function $\tau(x,y) : \Lambda^* \to \mathcal{H}$ by

$$(x,y) \mapsto (x,h(x,y))$$

for the hemishphere $h(x,y)$ containing x and $L(x,y)$. Note that $\tau(a) = \tau(a,r)$ if a is an attracting fixed point of an element of Γ with the repelling fixed point r in Λ.

We state a fact we need.

Lemma 19. *Let $\{(x_i,y_i) \in \Lambda^*\}$ be a sequence. Suppose that the segments $\{\overline{x_i y_i}\}$ form a sequence converging to the segment $\overline{xy}$ with $(x,y) \in \Lambda^*$ where $\{x_i\} \to x$ and $\{y_i\} \to y$. Then the corresponding sequence of invariant geodesics $L(x_i,y_i)$ converges to the line $L(x,y)$, and $\tau(x_i,y_i) \to \tau(x,y)$ as $i \to \infty$.*

Proof. This follows by the fact that $\tilde{N}$ is continuous by Proposition 11. □

□

Lemma 20. *Let $(x,y) \in \Lambda^*$. Then $\tau(x,y)$ does not depend on y, and $h(x,y)$ is never a hemisphere in $\mathbb{S}$ for every $(x,y) \in \Lambda^*$.*

Proof. Suppose that x is an endpoint of two weakly recurrent geodesics l with the starting point y and another one l' with the starting point y'.

If $(x, y) \in \Lambda^*$, then $L(x, y)$ is a line in $\mathsf{E}^{2,1}$ by Proposition 11; $h(x, y)$ is not in $\mathbb{S}$.

Choose a sequence $\{p_i\}$ of points $\overline{xy}^o$ so that $p_i \to x$. There exists a sequence $\{\gamma_i^{-1}\}$ of elements of Γ so that $\gamma^{-1}(p_i) \in F$ for a compact fundamental domain F of Ω_+ under Γ. (Recall that $\Sigma''_+ = \Omega_+/\Gamma$ is a compact hyperbolic surface with geodesic boundary and the weakly recurrent geodesics are always in it.)

By choosing a subsequence, we may assume that $\{p_i\} \to p_\infty \in F$ and $\{\gamma_i^{-1}(\mathrm{Cl}(l))\}$ converges to a segment s_∞ passing p_∞ and ending at two points of Λ. Let l_∞ denote the interior of s_∞ in Ω_+. Let x_∞ and y_∞ denote the endpoints so that we may assume without loss of generality that $\{\gamma_i^{-1}(x)\} \to x_\infty$ and $\{\gamma_i^{-1}(y)\} \to y_\infty$.

Since Γ is not elementary, we may assume that $\{x, y\} \cap \{x_\infty, y_\infty\} = \emptyset$ by multiplying γ_i^{-1} by an element of Γ if necessary.

Since $\{\gamma_i^{-1}(x)\} \to x_\infty$ and $\{\gamma_i^{-1}(y)\} \to y_\infty$, we have

$$\{\mathrm{Cl}(L(\gamma_i^{-1}(x), \gamma_i^{-1}(y))) = \gamma_i^{-1}(\mathrm{Cl}(L(x, y)))\} \to \mathrm{Cl}(L(x_\infty, y_\infty))$$

by continuity of $\mathscr{N}$ by Propositon 11. Hence.

$$\{\gamma_i^{-1}(h(x, y)) = h(\gamma_i^{-1}(x), \gamma_i^{-1}(x))\} \to h(x_\infty, y_\infty). \tag{20}$$

By using perpendicular geodesics in $\mathbb{S}_+$ to l, we can show that for an attracting fixed point a_i and repelling fixed point r_i of γ_i^{-1} for each i, it follows that $\{a_i\} \to y_\infty$ and $\{r_i\} \to x$. Since we have $\overline{r_i a_i} \to \overline{x y_\infty}$, it follows that

$$\begin{aligned} &(21) \qquad \{\mathrm{Cl}(L_{\gamma_i^{-1}}) = \mathrm{Cl}(L(r_i, a_i))\} \quad \to \quad \mathrm{Cl}(L(x, y_\infty)) \\ &(22) \qquad \{h(r_i, a_i)\} \quad \to \quad h(x, y_\infty). \end{aligned}$$

Since each a_i is an attracter, we obtain that $\gamma_i^{-1}(y') \to y_\infty$, and

$$\{\gamma_i^{-1}(h(x, y')) = h(\gamma_i^{-1}(x), \gamma_i^{-1}(y'))\} \to h(x_\infty, y_\infty). \tag{23}$$

Since $\partial h(x, y)$ is a great circle tangent to $\mathrm{bd}\mathbb{S}_+$ at x and so is $\partial h(x, y')$, we obtain $\partial h(x, y) = \partial h(x, y')$. We have that $\gamma_i^{-1}(\partial h(x, y'))$ is tangent to $\mathrm{bd}\mathbb{S}_+$ at $\gamma_i^{-1}(x)$. Thus, $\{\gamma_i^{-1}(\partial h(x, y'))\}$ converges to a great circle $\mathbf{S}^1_{x_\infty}$ tangent to $\mathrm{bd}\mathbb{S}_+$ at x_∞ in $\mathbb{S}$.

Choose a point q in the interior of the hemisphere corresponding to $h(x, y')$ bounded away from $\partial h(x, y) = \partial h(x, y')$.

Suppose that $q \in \mathscr{H} - h(x, y_\infty)$ for contradiction. Then q is uniformly bounded away from $h(r_i, a_i)$ for $i > I_0$ for a fixed integer I_0 by equation 21. We claim that $\{\gamma_i^{-1}(q)\} \to y_\infty$ or $\{\gamma_i^{-1}(q)\} \to y_{\infty,-}$:

We write γ_i^{-1} in the matrix of the form of equation 10 with respect to coordinates h_i so that $h_i \gamma_i^{-1} h_i^{-1}$ is of form

$$\hat{\gamma}_i^{-1} := \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \lambda(\gamma_i^{-1}) & 0 & 0 \\ k(\gamma_i^{-1}) & 0 & 1 & 0 \\ 0 & 0 & 0 & \frac{1}{\lambda(\gamma_i^{-1})} \end{bmatrix} \tag{24}$$

where $\lambda(\gamma_i^{-1}) > 1, k(\gamma_i^{-1}) > 0$ where

$$h_i(a_i) = [0,1,0,0], \qquad h_i(\hat{p}_i) = [1,0,0,0],$$
$$h_i(r_i) = [0,0,0,1], \qquad h_i(\hat{s}_i) = [0,0,1,0];$$

where we choose $\hat{p}_i \in L_{\gamma_i^{-1}}$ so that $\hat{p}_i$ is uniformly bounded away from $\mathbb{S}$ by equation 21, and $\hat{s}_i$ is the saddle-type fixed point or the point corresponding to an end point of $L_{\gamma_i^{-1}}$ in $\mathbb{S}$. As above, h_i is uniformly quasi-isometric with respect to $\mathbf{d}_{\mathbf{S}^3}$ and can be assume to be convergent. Also, $h(r_i, a_i)$ is the image of the hemisphere in $\mathscr{H}$ given by $\mathbf{S}_0^2$ of Lemma 12 under h_i^{-1}. (We are exactly in the situation of the part (I) of the proof of Proposition 13 in Section 4.) As before, we have $k(\gamma_i^{-1}) \leq C \log \lambda(\gamma_i^{-1})$ for the terms in the matrices of γ_i^{-1} by Lemma 14. By the first item of Lemma 12, we obtain $\{\gamma_i^{-1}(q)\} \to y_\infty$ or $\{\gamma_i^{-1}(q)\} \to y_{\infty,-}$ since h_i are uniformly quasi-isometric and q is bounded away from $h(r_i, a_i)$ uniformly for $i > I_0$.

Since $\{\gamma_i^{-1}(q)\} \to \pm y_\infty \in \mathbb{S}$ and $\{\gamma_i^{-1}(\partial h(x, y'))\}$ converges to a great circle $\mathbf{S}^1_{x_\infty}$ tangent to $\mathrm{bd}\mathbb{S}_+$ at x_∞ in $\mathbb{S}$, it follows that $\{\gamma_i^{-1}(h(x, y')\}$ converges to a hemisphere in $\mathbb{S}$ by Lemma 5. We obtained a contradiction to equation 23.

Therefore, $q \in h(x, y_\infty)$ and $h(x, y') = h(x, y_\infty)$ by geometry.

Similarly, we have $h(x, y) = h(x, y_\infty)$ by equation 20, and we conclude that $h(x, y) = h(x, y')$.

□ □

Hence, we have defined the function $\tau : \Lambda \to \mathcal{H}$ and the hemisphere $h(x)$ for any $x \in \Lambda$. Now, we show that τ is continuous.

Now suppose that $\{x_i\}$ is a sequence of arbitrary points in Λ so that $\{x_i\} \to x$. We choose $y_i \in \Lambda$ for each i so that $\{y_i\} \to y \in \Lambda$ for $y \neq x$. Then we have $\{h(x_i, y_i)\} \to h(x, y)$ by Lemma 19 and therefore, we have $\{\tau(x_i)\} \to \tau(x)$ by the independence of Lemma 20. Therefore, τ is a continuous function. □ □

Lemma 20 also generalizes the result in [8] that any weakly recurrent geodesics in $\mathsf{E}^{2,1}/\Gamma$ with the same forward limit set share the contracting subspace.

6.2. **Domains.** In $\mathbb{S}$, for each point x of Λ, we have a closed hemisphere $H_x \subset \mathbb{S}$ supporting the convex hull of Λ. We form an intersection $D(\Lambda) =$

$\bigcap_{x\in\Lambda} H_x^o \subset \mathbb{S}$. Then $D(\Lambda)$ is an open properly convex domain in $\mathbb{S}$. We note that

$$D(\Lambda) = \bigcup_{i\in\mathcal{J}} \triangle_i \cup \mathbb{S}_+ \tag{25}$$

where $\triangle_i$ is a $\mathbf{g}_i$-invariant triangle containing $\mathbf{l}_i$ as an edge where $\mathbf{g}_i$ is a corresponding element of Γ acting on $\mathbf{l}_i$ and hence its attracting and repelling fixed points a_i, r_i, at the ends and a saddle-type fixed point s_i are vertices of $\triangle_i$. Note that $\triangle_i \subset \mathcal{R}_i$ for the rectangular domain $\mathcal{R}_i$ defined in Section 2. Moreover, $D(\Lambda)/\Gamma$ is an open surface obtained from Σ_+ by adding open elementary annuli $\triangle_i/\langle\mathbf{g}_i\rangle$ in corresponding annuli A_j. Thus, the closure of $D(\Lambda)/\Gamma$ in Σ is a compact surface in Σ with boundary components $\mathbf{c}_j \subset A_j$ for each $j = 1, \dots, \mathsf{b}$.

Considering $D(\mathscr{A}(\Lambda)) = \mathscr{A}(D(\Lambda))$, we obtain closed curves $\mathbf{c}'_j \subset A_j$ for each $j = 1, \dots, \mathsf{b}$ that are boundary components of $D(\mathscr{A}(\Lambda))/\Gamma$ in Σ. Note that $(\Sigma - D(\Lambda) - D(\mathscr{A}(\Lambda)))/\Gamma$ is a union of compact annuli in each of A_j, $j = 1, \dots, \mathsf{b}$. We denote each component annulus by B_j so that $B_j \subset A_j$ where $\partial B_j = \mathbf{c}_j \cup \mathbf{c}'_j$. (Each B_j is a sum of two elementary annuli and is bounded by nonprincipal boundary components.)

The boundary of a convex open domain in $\mathsf{E}^{2,1}$ is *asymptotic* to a set K in $\mathbb{S}_+$ if every unbounded sequence of points in it has all limit points in K.

Corollary 21. *In $\mathsf{E}^{2,1}$, there exists a Γ-invariant nonempty convex open domain $\mathcal{D}$ whose boundary in $\mathsf{E}^{2,1}$ is asymptopic to* $\mathrm{bd}D(\Lambda)$*, homeomorphic to a circle. There exists another Γ-invariant convex open domain $\mathcal{D}'$ whose boundary in $\mathsf{E}^{2,1}$ is asymptotic to $\mathscr{A}(\mathrm{bd}D(\Lambda))$ so that the closures of $\mathcal{D}$ and $\mathcal{D}'$ are disjoint. Moreover, every weakly recurrent space-like geodesic is contained in a manifold*

$$(\mathsf{E}^{2,1} - \mathcal{D} - \mathcal{D}')/\Gamma$$

with concave boundary.

Proof. For each $x \in \Lambda$, take the open half-space $\mathcal{H}_x \subset \mathsf{E}^{2,1}$ bounded by $h(x, y)$ for some $y \in \Lambda - \{x\}$ and containing Λ in its closure in the hemisphere $\mathscr{H}$. Then we take $\mathcal{D} := \bigcap_{x\in\Lambda} \mathcal{H}_x$, which is a convex open domain.

We show that $\mathcal{D}$ is not empty: That is, we claim that there exists a compact convex set $K \subset \mathsf{E}^{2,1}$ so that $\mathcal{H}_x \cap K \neq \emptyset$ for all $x \in \Lambda$ and hence there exist

$$z \in K \cap \bigcap_{x\in\Lambda} \mathrm{Cl}_{\mathbf{S}^3}(\mathcal{H}_x)$$

by compactness. Then the join $J(z, \Lambda)$ is a subset of $\mathrm{Cl}_{\mathbf{S}^3}(\mathcal{H}_x)$ for every $x \in \Lambda$. Hence the topological interior of the join $J(z, \Lambda)$ is a nonempty subset of $\mathcal{D}$: Suppose not. Then there exists a sequence of hemispheres $h(x_i, y_i), x_i, y_i \in \Lambda$ so that it bounds an open half-space $\mathcal{H}_{x_i}$ in $\mathsf{E}^{2,1}$ and $\{\mathcal{H}_{x_i} \cap K\}$ for each compact set $K \subset \mathsf{E}^{2,1}$ is eventually empty as $i \to \infty$. Since Λ is compact, this is equivalent to saying that $\{h(x_i, y_i)\}$ converges to a hemisphere in $\mathbb{S}$ up to a choice of a subsequence. Let x_∞ be a limit point of the sequence $\{x_i\}$. Then $\tau(x_\infty)$ is a pair (x_∞, h_∞) for a hemisphere h_∞ in $\mathbb{S}$ by the continuity of τ contradicting Lemma 20.

Let $\mathrm{bd}_{\mathsf{E}^{2,1}}(\bigcap_{x\in\Lambda}\mathcal{H}_x)$ be the topological boundary in $\mathsf{E}^{2,1}$. The closure of $\bigcap_{x\in\Lambda}\mathcal{H}_x$ is a convex compact subset of $\mathbf{S}^3$, and the boundary of $\bigcap_{x\in\Lambda}\mathcal{H}_x$ in $\mathbf{S}^3$ equals $\mathrm{Cl}(D(\Lambda)) \cup \mathrm{bd}_{\mathsf{E}^{2,1}}(\bigcap_{x\in\Lambda}\mathcal{H}_x)$. Thus, every unbounded sequence of points in $\mathrm{bd}_{\mathsf{E}^{2,1}}(\bigcap_{x\in\Lambda}\mathcal{H}_x)$ will have limit points in $D(\Lambda)$. Since the limit points cannot be in $D(\Lambda)^o$, we see that they are in $\mathrm{bd}D(\Lambda)$.

For each $\mathcal{H}_x$, we can take $(\mathsf{E}^{2,1} - \mathcal{H}_x)^o$ and take the intersection over all $x \in \Lambda$. This gives us $\mathcal{D}'$ with the right property.

First, notice $\mathcal{D} \cap \mathcal{D}' = \emptyset$. If $\mathrm{Cl}(\mathcal{D}) \cap \mathrm{Cl}(\mathcal{D}') \neq \emptyset$, then it is a compact convex subset of $\mathsf{E}^{2,1}$ by the asymptotic properties and the fact that

$$\mathrm{Cl}(D(\Lambda)) \cap \mathrm{Cl}(\mathscr{A}(D(\Lambda))) = \emptyset.$$

Since $\mathrm{Cl}(\mathcal{D}) \cap \mathrm{Cl}(\mathcal{D}')$ is also Γ-invariant, the Brouwer fixed-point theorem implies a contradiction that the intersection contains a fixed point for each element $g \in \Gamma$. Hence, $\mathrm{Cl}(\mathcal{D}) \cap \mathrm{Cl}(\mathcal{D}') = \emptyset$.

Every weakly recurrent space-like geodesics equals $L(x, y)$ for $(x, y) \in \Lambda^*$ by Theorem 1 of [27]. Since $L(x, y)$ is in $h(x, y)$ for some $(x, y) \in \Lambda^*$, the last statement follows. $\square$ $\square$

Remark 22. These domains are related to the invariant domains found by Mess [33] for the closed surface groups.

7. GEOMETRIC FINITENESS

We define geometric finiteness and show that the Margulis space-times are geometrically finite.

We start with

Theorem 23 (Scott [36])**.** *If M is an orientable 3-manifold and $\pi_1(M)$ is finitely generated, then there exists a compact submanifold N of M such that the inclusion homomorphism induces an isomorphism $\pi_1(N) \to \pi(M)$.*

If M is aspherical, as in Margulis space-times, then we can construct a compact submanifold so that the inclusion map $N \to M$ is a homotopy equivalence. We say that a compact submanifold with properties of N is a *core* of M. For $\mathsf{E}^{2,1}/\Gamma$ with a free group Γ of a finite rank, there exists a core.

Definition 24 (Geometric finitness)**.** Let Γ be the orientation-preserving Lorentzian isometry group acting on the Lorenzian space $\mathsf{E}^{2,1}$ freely and properly discontinuously. Let $\mathsf{E}^{2,1}/\Gamma$ be given an arbitrary complete Riemannian metric δ and a core M_0. $\mathsf{E}^{2,1}/\Gamma$ is *geometrically finite* if for each pair of a closed geodesic c and a point x on c in $\mathsf{E}^{2,1}/\Gamma$, x is of a uniformly bounded δ-distance from M_0; that is, $\delta(x, M_0) \leq C$ for a constant C independent of x and c.

Actually, since M_0 is compact, we can replace M_0 by a base point x_0 in the above definition.

The geometrical finiteness here of course is already discovered by Goldman, Labourie, and Margulis [28], [27], and we only restate it in this form.

Lemma 25. *Fix an arbitrary point x_0 and an arbitrary Euclidean metric d_E on $\mathsf{E}^{2,1}$ seen as an affine space. A Margulis space-time $\mathsf{E}^{2,1}/\Gamma$ is geometrically finite if and only if there exists a constant $C' > 0$ such that for any $g \in \Gamma$ acting on a complete space-like line L_g where it acts on as a translation and a point $t \in L_g$ in $\mathsf{E}^{2,1}$, there exists $\gamma \in \Gamma$ so that $\gamma(t)$ is of a uniformly bounded distance from x_0 by C' with respect to the Euclidean metric d_E.*

Proof. Let $C > 0$ be the uniform bound for the Riemannian metric of $\mathsf{E}^{2,1}/\Gamma$ used in Definition 24. A δ-ball of radius C in $\mathsf{E}^{2,1}$ maps to a bounded ball in $\mathsf{E}^{2,1}$ in the Euclidean metric d_E since it is compact. The Euclidean diameters of the δ-balls of radii equal to C with centers in a compact set K are bounded above by a constant $C' > 0$.

Given $g \in \Gamma$ acting on L_g and $t \in L_g$, we have that L_g maps to a closed geodesic c and a point $x \in c$ from a point $x_0 \in M_0$ of the δ-distance bounded above by C. Let F be the compact fundamental domain of M_0 in $\mathsf{E}^{2,1}$. Assume without loss of generality that $x_0 \in F$. Choose $\gamma \in \Gamma$ so that $\gamma(t)$ is of the δ-distance $\leq C'$ from $x' \in F$. Thus, $\gamma(t)$ is of the Euclidean distance $\leq C''$ from x_0 for some $C'' > 0$.

Conversely, a Euclidean ball with center x_0 of radius C' has a δ-diameter uniformly bounded above. The proof of the converse is very similar to the above. □ □

Proposition 26. *Let Γ be a nonamenable orientation-preserving Lorentzian isometry group acting on the Lorenzian space $\mathsf{E}^{2,1}$ freely and properly discontinuously. Then the Margulis space-time $\mathsf{E}^{2,1}/\Gamma$ is geometrically finite.*

Proof. Consider the compact fundamental domain $\hat{F}$ of $\mathbb{U}_{\mathrm{rec}}\mathbb{S}_+ = \widehat{\mathbb{U}_{\mathrm{rec}}\Sigma_+}$ under the deck transformation group of covering to $\mathbb{U}_{\mathrm{rec}}\Sigma_+$. This is a closed subset of the compact set $\mathbb{U}\Sigma''_+$. Then $\pi_{\mathsf{E}^{2,1}} \circ \tilde{N}(\hat{F})$ gives us a set of geodesics of bounded distances from x_0 since it is an image of a compact set.

Let g be an element of Γ, and let t be a point on the line L_g in $\mathsf{E}^{2,1}$ where g acts as a translation. Let u denote the unit vector at t tangent to L_g. Then (t, u) correspond to (t', u') for a point $t' \in \mathbb{S}_+$ and u' on $\mathbb{U}_{\mathrm{rec}}\mathbb{S}_+$ via $\pi_{\mathsf{E}^{2,1}} \circ \tilde{N}$ by Proposition 11. Since $\hat{F}$ is a fundamental domain, $(\gamma(t'), D\gamma(\mathsf{u}'))$ is in the bounded set $\hat{F}$ for a deck transformation group γ. (Here by $D\gamma : \mathbb{U}\mathbb{S}_+ \to \mathbb{U}\mathbb{S}_+$, we mean the differential map.) □ □

Theorem 27. *There exists a core in a Margulis space-time containing all weakly recurrent space-like geodesics.*

Proof. Let Γ be a free group of rank ≥ 2 acting freely and properly discontinuously on $\mathsf{E}^{2,1}$. We construct $\tilde{\Sigma}$ as above.

Let M denote the compact handlebody $(\mathsf{E}^{2,1} \cup \tilde{\Sigma})/\Gamma$ so that M^o is the Margulis space-time $\mathsf{E}^{2,1}/\Gamma$. We denote by $\tilde{M}$ the space $\mathsf{E}^{2,1} \cup \tilde{\Sigma}$. Recall the domains $\mathcal{D}$ and $\mathcal{D}'$ from the proof of Corollary 21. Then consider the closure $\hat{\mathcal{D}} = \mathrm{Cl}_{\tilde{M}}(\mathcal{D})$ of $\mathcal{D}$ in $\tilde{M}$. Then Γ acts properly discontinuously on $\hat{\mathcal{D}}$ as well.

The topological boundary $\mathrm{bd}_{\tilde{M}}\hat{\mathcal{D}}$ in $\tilde{M}$ is a closed invariant subset of $\mathsf{E}^{2,1}$ and is a connected properly imbedded 2-dimensional submanifold invariant under Γ. Since $\mathrm{bd}_{\tilde{M}}\hat{\mathcal{D}}$ equals $\partial\mathrm{Cl}(\mathcal{D}) - D(\Lambda)$, $\mathrm{bd}_{\tilde{M}}\hat{\mathcal{D}}$ is homeomorphic to a 2-cell. However, under projection $p : \tilde{M} \to M$, it goes to a compact closed connected properly imbedded surface Σ_1 bounding the image of $\hat{\mathcal{D}}$. To explain more, $M_1 := \hat{\mathcal{D}}/\Gamma$ is imbedded in M and is homotopy equivalent to M and the topological boundary $\mathrm{bd}_M M_1$ in M is the compact surface Σ_1.

Since Σ_1 is homotopic to the closure S_1 of $D(\Lambda)/\Gamma$ in Σ sharing boundary components in M_1 and $\partial M_1 = \Sigma_1 \cup S_1$, it follows that M_1 is homeomorphic to $\Sigma_1 \times I$ for an interval I by Theorem 10.2 of [29].

Notice also

$$\partial\Sigma_1 = \partial S_1 = \bigcup_{i=1}^{\mathsf{b}} \mathbf{c}_i$$

for the number of boundary components b of S_1.

We can do similarly for the closure $\hat{\mathcal{D}}_- := \mathrm{Cl}_{\tilde{M}}(\mathcal{A}(\mathcal{D}))$ and we obtain $M_2 := \hat{\mathcal{D}}_-/\Gamma$ imbedded into M disjoint from M_1. Similary M_2 is homeomorphic to $\Sigma_2 \times I$ for a surface $\Sigma_2 = \mathrm{bd}_M M_2$ which is also homotopy equivalent to M.

Notice also

$$\partial\Sigma_2 = \bigcup_{i=1}^{\mathsf{b}} \mathbf{c}'_i.$$

Let $\mathrm{int}M_1$ and $\mathrm{int}M_2$ denote the topological interiors. Since we are removing the product of I with surfaces, it follows that $M - \mathrm{int}M_1 - \mathrm{int}M_2$ is a compact 3-manifold homeomorphic to M bounded by $\Sigma_1 \cup \Sigma_2 \cup B_1 \cup \cdots B_g$. (Recall B_i from Section 6.2.)

By geometric finiteness, every closed geodesic is of a uniformly bounded δ-distance from a base point x_0 in M. Therefore, we can isotopy all B_i to imbedded annuli in M^o so that the resulting imbedded annuli B'_i are all disjoint from closed space-like geodesics in M^o and $\partial B'_i \subset \Sigma_1 \cup \Sigma_2$. Therefore, Σ_1 and Σ_2 and $B'_1, \ldots, B'_g$ bound a compact submanifold $\hat{M}$ containing all closed space-like geodesics. Since by Anosov property of the space-like geodesic flow in M^o, the closed space-like geodesics are dense in the space of weakly recurrent geodesics by Theorem 1 of [27], $\hat{M}$ is the desired compact submanifold in M^o. □ □

References

[1] Anderson, J., Bonfert-Taylor, P., Taylor, E.: Convergence groups, Hausdorff dimension, and a theorem of Sullivan and Tukia. Geom. Dedicata **103**, 51–67 (2004)

[2] Abels, H., Margulis, G.A., Soifer, G.A.: Properly discontinuous groups of affine transformations with orthogonal linear part. C. R. Acad. Sci. Paris Sér. I Math. **324**(3), 253–258 (1997)

[3] Beardon, A.: Geometry of discrete groups. Graduate Text in Mathematics Vol. 91, Springer New York, 1995

[4] Benoist, Y., Convexes divisibles I In: Dani, S.G., Prasad, G. (eds.) Algebraic groups and arithmetic, Tata Inst. Fund. Res. Stud. Math. 17, pp. 339–374 Tata Institute of Fund. Research (2004)

[5] Bowditch, B., Notes on tameness. Enseign. Math. (2) **56**(3-4), 229–285 (2010)

[6] Browder, W., Levine, J., Livesay, G.R.: Finding a boundary for an open manifold. Amer. J. Math. **87**, 1017–1028 (1965)

[7] Charette, V., Drumm, T., Goldman, W.: Affine deformations of the three-holed sphere. Geometry & Topology **14**, 1355–1382 (2010)

[8] Charette, V., Goldman, W., Jones, C.: Recurrent geodesics in flat Lorentzian manifolds. Canadian Math. Bull. **47**(3), 332–342 (2004)

[9] Choi, S., The Margulis Lemma and the thick and thin decomposition for convex real projective surfaces. Advances in mathematics **122**, 150–191 (1996)

[10] Choi, S., Convex decompositions of real projective surfaces. I: *π-annuli and convexity.* J. Differential Geom. **40**(1), 165–208 (1994)

[11] Choi, S., Convex decompositions of real projective surfaces. II: *Admissible decompositions.* J. Differential Geom. **40**(2), 239–283 (1994)

[12] Choi, S., Goldman, W., The classification of real projective structures on compact surfaces. Bull. Amer. Math. Soc. (N.S.) **34**(2), 161–171 (1997)

[13] Cooper, D., Long, D., Thistlethwaite, M.: Computing varieties of representations of hyperbolic 3-manifolds into $\mathrm{SL}(4, \mathbb{R})$. Experiment. Math. **15**, 291–305 (2006)

[14] D. Cooper, D., Long, D., Thistlethwaite, M.: Flexing closed hyperbolic manifolds. Geom. Topol. **11**, 2413–2440 (2007)

[15] Drumm, T., Fundamental polyhedra for Margulis space-times. Doctoral dissertation, Unversity of Maryland (1990); Topology **31**(4), 677–683 (1992)

[16] Drumm, T., Linear holonomy of Margulis space-times. J. Differential Geometry **38**, 679–691 (1993)

[17] Drumm, T., Goldman, W.: The Geometry of Crooked Planes. Topology **38**(2), 323–351 (1999)

[18] Drumm, T., Goldman, W.: The isospectrality of flat Lorentz 3-manifolds. J. Differential Geometry **58**, 457–465 (2001)

[19] Eberlein, P.: Geodesic flows on negatively curved manifolds: I. Annals of Mathematics, 2nd Ser. **95**, 492–510 (1972)

[20] Francis, C: The conformal boundary of Margulis space-times. C.R. Acad. Sci. Paris, Ser. **1336**, 751–756 (2003)

[21] Fried, D., Goldman, W., Hirsch, M.: Affine manifolds with nilpotent holonomy. Comment. Math. Helv. **56**(4), 487–523 (1981)

[22] Fried, D., Goldman, W.: Three-dimensional affine crystallographic groups. Adv. in Mathematics **47**(1), 1–49 (1983)

[23] Goldman, W: Affine manifolds and projective geometry on surfaces. Senior Thesis, Princeton University, 1977

[24] Goldman, W.: Projective structures with Fuchsian holonomy. J. Differential Geom. **25**(3), 297–326 (1987)

[25] Goldman, W.: Convex real projective structures on compact surfaces. J. Differential Geometry **31**, 791–845 (1990)

[26] Goldman, W.: Projective geometry of manifolds. Lecture notes from a 1988 course. `http://www2.math.umd.edu/~wmg/pgom.pdf` (1988)

[27] Goldman, W., Labourie, F.: Geodesics in Margulis space times. Ergod. Th. & Dynamic. Sys. **32**, 643–651 (2012)

[28] Goldman, W., Labourie, F., Margulis, G.: Proper affine actions and geodesic flows of hyperbolic surfaces, Annals of Mathematics **170**, 1051–1083 (2009)

[29] J. Hempel, 3-manifolds Princeton University Press, Princeton, 1976.

[30] Katok, S., Hasselblatt, B.: Introduction to the modern theory of dynamical systems. Encyclopedia of mathematics and its applications vol. 54, Cambridge University Press, Cambridge 1995

[31] Marden, A.: Outer circles: an introduction to hyperbolic 3-manifolds. Cambridge University Press, Cambridge, 2007

[32] Margulis, G. A.: Complete affine locally flat manifolds with a free fundamental group. In: Automorphic functions and number theory, II. Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) 134, 190–205 (1984)

[33] Mess, G.: Lorentz space-times of constant curvature. Geom. Dedicata **126**, 3–45 (2007)

[34] Meyers, R.: End reductions, fundamental groups, and covering spaces of irreducible open 3-manifolds. Geometry & Topology **9**, 971–990 (2005)

[35] Nagano, T.,Yagi, K.: The affine structures on the real two-torus: I. Osaka J. of Math. **11**, 181–210 (1974)

[36] Scott, P.: Compact submanifolds of 3-manifolds. J. London Math. Soc. (2) **7**, 246–250 (1973)

[37] Siebenmann, L.C.: The obstruction to finding a boundary for an open manifold of dimension greater than five. Ph.D. thesis, Princeton University (1965)

[38] Thurston, W.: Geometry and topology of 3-manifolds. Lecture notes. Princeton University `http://www.msri.org/publications/books/gt3m/` (1979)

[39] Scott, P., Tucker, T.: Some examples of exotic noncompact 3-manifolds. Quart. J. Math. Oxford Ser. (2) **40**(160), 481–499 (1989)

[40] Tucker, T.: Non-compact 3-manifolds and the missing-boundary problem. Topology **13**, 267–273 (1974)

Department of Mathematical Sciences, KAIST, Daejeon 305-701, South Korea

E-mail address: `schoi@math.kaist.ac.kr`